


Crystal structure of [1,2-ethylene-1,1'-bis(η^5 -tetrahydroindenyl)]-chloro- $[\eta^2$ -N,C-3,4,5,6-tetrafluoropyridyl]zirconium(IV), $\text{ZrCl}(\text{C}_5\text{F}_4\text{N})(\text{C}_{20}\text{H}_{24})$

A. Spannenberg*, U. Jäger-Fiedler, P. Arndt and U. Rosenthal

Leibniz-Institut für Organische Katalyse an der Universität Rostock e.V., Buchbinderstr. 5–6, 18055 Rostock, Germany

Received March 21, 2005, accepted and available on-line April 8, 2005; CCDC no. 1267/1513



Abstract

$\text{C}_{25}\text{H}_{24}\text{ClF}_4\text{NZr}$, monoclinic, $P12_1/n1$ (no. 14), $a = 9.903(2) \text{ \AA}$, $b = 11.279(2) \text{ \AA}$, $c = 20.231(4) \text{ \AA}$, $\beta = 92.43(3)^\circ$, $V = 2257.7 \text{ \AA}^3$, $Z = 4$, $R_{\text{gt}}(F) = 0.035$, $wR_{\text{obs}}(F^2) = 0.076$, $T = 200 \text{ K}$.

Source of material

The complex $\text{rac}-(\text{ebthi})\text{Zr}(\eta^2\text{-Me}_3\text{SiC}_2\text{SiMe}_3)$ (1.0 g, 1.9 mmol; ebthi = 1,2-ethylene-1,1'-bis(tetrahydroindenyl)) was dissolved in 10 ml of toluene under Ar and treated under stirring with pentafluoropyridine (0.8 ml, 7.3 mmol). Traces of 2-chloro-3,4,5,6-tetrafluoropyridine contained in the batch of pentafluoropyridine reacted with $\text{rac}-(\text{ebthi})\text{Zr}(\eta^2\text{-Me}_3\text{SiC}_2\text{SiMe}_3)$. After 15 min at 353 K the green solution became green-brownish and a light brown precipitate had formed. The solvent was removed in vacuum, and the residue was dissolved in $[\text{D}_6]$ -benzene for NMR measurements. From this solution a few beige crystals were obtained.

Experimental details

Both anellated cyclohexyl rings of the ebthi ligand are disordered. The corresponding atoms were refined using split positions.

Discussion

The activation of several types of C–F bonds by many transition metal complexes was summarized in many reviews [1–6]. Beck-

haus and coworkers observed the C–F bond cleavage reaction of pentafluoropyridine with $\text{Cp}_2\text{Ti}(\eta^2\text{-Me}_3\text{SiC}_2\text{SiMe}_3)$ under formation of a F-bridged dinuclear Ti(III) complex [7]. In the corresponding reaction with $\text{Cp}_2\text{Zr}(L)(\eta^2\text{-Me}_3\text{SiC}_2\text{SiMe}_3)$ (L = pyridine), a C–F activation took place and the formed $\text{Cp}_2\text{Zr}(4\text{-C}_5\text{NF}_4)\text{F}$ was confirmed by crystal structure analysis [8].

The crystal structure of $\text{rac}-(\text{ebthi})\text{Zr}(\eta^2\text{-N,C-3,4,5,6-NC}_5\text{F}_4)\text{Cl}$, reveals C–Cl activation in 2-position instead of C–F activation of the 2-chloro-3,4,5,6-tetrafluoropyridine. The molecular structure of $\text{rac}-(\text{ebthi})\text{Zr}(\eta^2\text{-N,C-3,4,5,6-NC}_5\text{F}_4)\text{Cl}$ ($d(\text{Zr1}—\text{Cl1}) = 2.241(3) \text{ \AA}$, $d(\text{Zr1}—\text{N1}) = 2.289(3) \text{ \AA}$, $d(\text{Cl1}—\text{N1}) = 1.348(5) \text{ \AA}$, $d(\text{Zr1}—\text{C11}) = 2.519(1) \text{ \AA}$, $\angle \text{N1}—\text{Zr}—\text{Cl1} = 34.6(1)^\circ$) is comparable with the cationic zirconocene complex $[\text{Cp}_2\text{Zr}(\eta^2\text{-N,C-picolyl})(\text{PMe}_3)]^+$ ($d(\text{Zr}—\text{C1}) = 2.29(2) \text{ \AA}$, $d(\text{Zr}—\text{N}) = 2.21(1) \text{ \AA}$, $d(\text{Cl}—\text{N}) = 1.32(2) \text{ \AA}$, $\angle \text{N}—\text{Zr}—\text{Cl} = 34.2(2)^\circ$) [9]. The Zr—Cl bond distance in the title compound is longer than that found for the corresponding $\text{rac}-(\text{ebthi})\text{ZrCl}_2$ [10], but similar to the Zr—Cl bond length in $[\text{SiMe}_2(\text{C}_5\text{H}_4)_2][(\eta^5\text{-C}_5\text{H}_5)\text{ZrCl}(\mu\text{-H})_2]$ ($d(\text{Zr1}—\text{Cl1}) = 2.526(1) \text{ \AA}$, $d(\text{Zr2}—\text{Cl2}) = 2.521(1) \text{ \AA}$) [11].

Table 1. Data collection and handling.

Crystal:	beige prism, size $0.2 \times 0.3 \times 0.4 \text{ mm}$
Wavelength:	Mo K_α radiation ($0.71073 \text{ \AA}$)
μ :	6.51 cm^{-1}
Diffraction, scan mode:	STOE IPDS I, φ
$2\theta_{\text{max}}$:	48.48°
$N(hkl)_{\text{measured}}$, $N(hkl)_{\text{unique}}$:	6533, 3561
Criterion for I_{obs} , $N(hkl)_{\text{gt}}$:	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 2716
$N(\text{param})_{\text{refined}}$:	285
Programs:	SHELXS-86 [12], SHELXL-97 [13]

Table 2. Atomic coordinates and displacement parameters (in $\text{\AA}^2$).

Atom	Site	Occ.	x	y	z	U_{iso}
H(6)	4e		0.9427	0.3627	0.7328	0.049
H(7)	4e		1.1826	0.2874	0.7346	0.049
H(11A)	4e		0.7494	0.3509	0.6145	0.071
H(11B)	4e		0.6979	0.2500	0.6632	0.071
C(12A)	4e	0.73(1)	0.7249(5)	0.1870(6)	0.5678(3)	0.061(2)
H(12A)	4e	0.73	0.6922	0.1099	0.5842	0.074
H(12B)	4e	0.73	0.6496	0.2241	0.5416	0.074
C(13A)	4e	0.73	0.8416(5)	0.1636(6)	0.5219(3)	0.061(2)
H(13A)	4e	0.73	0.8093	0.1151	0.4836	0.073
H(13B)	4e	0.73	0.8764	0.2396	0.5050	0.073
C(12B)	4e	0.27	0.744(2)	0.242(1)	0.5531(3)	0.056(5)
H(12C)	4e	0.27	0.7920	0.3023	0.5275	0.068
H(12D)	4e	0.27	0.6467	0.2433	0.5387	0.068
C(13B)	4e	0.27	0.8041(7)	0.118(1)	0.543(1)	0.067(6)

* Correspondence author (e-mail: anke.spannenberg@ifok-rostock.de)

Table 2. Continued.

Atom	Site	Occ.	x	y	z	U _{iso}
H(13C)	4e	0.27	0.7894	0.0980	0.4953	0.081
H(13D)	4e	0.27	0.7511	0.0612	0.5682	0.081
H(14A)	4e		0.9220	0.0182	0.5741	0.064
H(14B)	4e		1.0334	0.0873	0.5336	0.064
H(15)	4e		0.9693	-0.1768	0.7645	0.046
H(16)	4e		1.1014	-0.1044	0.6679	0.049
H(20A)	4e		1.0194	-0.1613	0.9004	0.062
H(20B)	4e		0.9867	-0.0227	0.9069	0.062
C(21A)	4e	0.55(1)	1.1815(6)	-0.0651(6)	0.9401(4)	0.059(3)
H(21A)	4e	0.55	1.2495	-0.1266	0.9311	0.070
H(21B)	4e	0.55	1.1577	-0.0710	0.9870	0.070
C(22A)	4e	0.55	1.2362(9)	0.0575(6)	0.9254(3)	0.057(3)
H(22A)	4e	0.55	1.3111	0.0749	0.9578	0.068

Table 2. Continued.

Atom	Site	Occ.	x	y	z	U _{iso}
H(22B)	4e	0.55	1.1638	0.1162	0.9323	0.068
C(21B)	4e	0.45	1.1466(8)	-0.005(1)	0.9404(5)	0.068(4)
H(21C)	4e	0.45	1.1052	0.0749	0.9428	0.082
H(21D)	4e	0.45	1.1462	-0.0397	0.9853	0.082
C(22B)	4e	0.45	1.2929(9)	0.011(1)	0.9223(4)	0.080(4)
H(22C)	4e	0.45	1.3433	0.0581	0.9564	0.096
H(22D)	4e	0.45	1.3377	-0.0672	0.9182	0.096
H(23A)	4e		1.2893	0.1611	0.8460	0.070
H(23B)	4e		1.3807	0.0444	0.8546	0.070
H(24A)	4e		1.3612	0.1269	0.7244	0.066
H(24B)	4e		1.3884	0.0065	0.6850	0.066
H(25A)	4e		1.2275	0.0512	0.6043	0.063
H(25B)	4e		1.3133	0.1706	0.6153	0.063

Table 3. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	x	y	z	U ₁₁	U ₂₂	U ₃₃	U ₁₂	U ₁₃	U ₂₃
Zr(1)	4e	0.98432(3)	0.09263(3)	0.74117(2)	0.0293(2)	0.0286(2)	0.0293(2)	-0.0012(2)	0.0027(1)	-0.0022(2)
Cl(1)	4e	0.7775(1)	-0.0151(1)	0.69600(5)	0.0529(6)	0.0627(7)	0.0576(7)	-0.0214(5)	-0.0135(5)	0.0028(5)
C(1)	4e	0.9641(4)	0.1951(3)	0.8351(2)	0.041(2)	0.043(2)	0.028(2)	0.009(2)	-0.000(2)	0.000(2)
C(2)	4e	0.9748(5)	0.2742(4)	0.8868(2)	0.084(3)	0.043(2)	0.039(2)	0.019(2)	-0.006(2)	-0.001(2)
C(3)	4e	0.8680(7)	0.2955(4)	0.9255(2)	0.129(5)	0.064(3)	0.035(2)	0.044(3)	0.017(3)	-0.002(2)
C(4)	4e	0.7494(6)	0.2401(5)	0.9126(3)	0.103(4)	0.090(4)	0.065(4)	0.047(4)	0.053(4)	0.012(3)
C(5)	4e	0.7398(5)	0.1627(5)	0.8601(3)	0.054(3)	0.081(4)	0.082(4)	0.011(3)	0.027(3)	0.013(3)
C(6)	4e	0.9774(4)	0.3049(3)	0.7039(2)	0.055(2)	0.024(2)	0.043(2)	0.003(2)	0.008(2)	0.000(2)
C(7)	4e	1.1116(4)	0.2635(3)	0.7045(2)	0.045(2)	0.031(2)	0.047(2)	-0.009(2)	0.003(2)	0.002(2)
C(8)	4e	1.1231(4)	0.1808(3)	0.6530(2)	0.045(2)	0.031(2)	0.047(2)	-0.001(2)	0.018(2)	0.009(2)
C(9)	4e	0.9941(4)	0.1682(3)	0.6223(2)	0.053(2)	0.029(2)	0.029(2)	0.005(2)	0.008(2)	0.003(2)
C(10)	4e	0.9042(4)	0.2461(3)	0.6531(2)	0.049(2)	0.034(2)	0.036(2)	0.008(2)	0.003(2)	0.005(2)
C(11)	4e	0.7606(4)	0.2667(4)	0.6276(2)	0.059(3)	0.063(3)	0.056(3)	0.025(2)	-0.007(2)	-0.004(2)
C(14)	4e	0.9542(4)	0.0977(4)	0.5613(2)	0.069(3)	0.056(2)	0.035(2)	0.008(2)	0.006(2)	-0.006(2)
C(15)	4e	1.0407(4)	-0.1205(3)	0.7667(2)	0.040(2)	0.025(2)	0.051(2)	-0.001(2)	0.001(2)	0.002(2)
C(16)	4e	1.1153(4)	-0.0807(3)	0.7127(2)	0.051(2)	0.034(2)	0.039(2)	0.013(2)	0.008(2)	-0.003(2)
C(17)	4e	1.2140(3)	0.0004(3)	0.7372(2)	0.033(2)	0.028(2)	0.056(2)	0.006(2)	0.007(2)	0.006(2)
C(18)	4e	1.1976(3)	0.0122(3)	0.8054(2)	0.033(2)	0.032(2)	0.047(2)	0.007(2)	-0.005(2)	-0.001(2)
C(19)	4e	1.0904(3)	-0.0625(3)	0.8236(2)	0.039(2)	0.029(2)	0.039(2)	0.004(2)	-0.001(2)	0.004(2)
C(20)	4e	1.0565(4)	-0.0806(4)	0.8948(2)	0.062(2)	0.052(2)	0.040(2)	0.005(2)	0.003(2)	0.005(2)
C(23)	4e	1.2873(4)	0.0753(4)	0.8563(2)	0.038(2)	0.050(3)	0.086(3)	0.003(2)	-0.015(2)	-0.006(2)
C(24)	4e	1.3176(4)	0.0632(4)	0.6974(2)	0.041(2)	0.050(3)	0.075(3)	0.008(2)	0.022(2)	0.009(2)
C(25)	4e	1.2503(4)	0.1153(3)	0.6363(2)	0.045(2)	0.049(3)	0.066(3)	0.007(2)	0.027(2)	0.014(2)
N(1)	4e	0.8439(3)	0.1410(3)	0.8241(2)	0.043(2)	0.053(2)	0.048(2)	0.008(2)	0.015(2)	-0.002(2)
F(1)	4e	1.0920(3)	0.3329(2)	0.9001(1)	0.114(2)	0.065(2)	0.070(2)	0.001(2)	-0.032(2)	-0.024(1)
F(2)	4e	0.8788(4)	0.3726(3)	0.9759(1)	0.233(4)	0.103(2)	0.047(2)	0.065(3)	0.018(2)	-0.026(2)
F(3)	4e	0.6410(4)	0.2596(4)	0.9491(2)	0.163(3)	0.157(4)	0.133(3)	0.068(3)	0.112(3)	0.016(3)
F(4)	4e	0.6256(3)	0.1064(4)	0.8446(2)	0.055(2)	0.153(3)	0.184(4)	-0.011(2)	0.059(2)	-0.011(3)

Acknowledgment. The work was supported by the Deutsche Forschungsgemeinschaft (project code SPP 1118).

References

- Kiplinger, J. L.; Richmond, T. G.; Osterberg, C. E.: Activation of Carbon-Fluorine Bonds by Metal Complexes. *Chem. Rev.* **94** (1994) 373-431.
- Burdeniuc, J.; Jedlicka, B.; Crabtree, R. H.: Recent Advances in C-F Bond Activation. *Chem. Ber./Recueil* **130** (1997) 145-154.
- Richmond, T. G.: Metallorganische Umsetzungen belegen die Reaktivität von Fluorkohlenwasserstoffen. *Angew. Chem.* **112** (2000) 3378-3380.
- Richmond, T. G.: Metal Reagents for Activation and Functionalization of Carbon-Fluorine Bonds. In: *Topics in Organometallic Chemistry*, Vol. 3 (Ed. S. Murai), p. 243-269. Springer, New York 1999.
- Jones, W. D.: Activation of C-F bonds using Cp*₂ZrH₂: a diversity of mechanisms. *Dalton Trans.* (2003) 3991-3995.
- Mazurek, U.; Schwarz, H.: Carbon fluorine bond activation looking at and learning from unsolvated systems. *Chem. Commun.* (2003) 1321-1326.
- Piglosiewicz, I. M.; Kraft, S.; Beckhaus, R.; Haase, D.; Saak, W.: Selective C-H and C-F Bond Activation Reactions of Pyridine and Fluoropyridines—

- Formation of Binuclear μ -X Titanocene Complexes (X = H, F) with α -Functionalized N-Heterocycles. *Eur. J. Inorg. Chem.* (2005) 938-945.
- Jäger-Fiedler, U.; Arndt, P.; Baumann, W.; Spannenberg, A.; Burlakov, V. V.; Rosenthal, U.: Reactions of Zirconocene Bis(trimethylsilyl)acetylene Complexes with Fluorinated Pyridines: C-H versus C-F Bond Activation. *Eur. J. Inorg. Chem.* (2005), in press.
 - Jordan, R. F.; Taylor, D. F.; Baenziger, N. C.: Synthesis and Insertion Chemistry of Cationic Zirconium(IV) Pyridyl Complexes. Productive σ -Bond Metathesis. *Organometallics* **9** (1990) 1546-1557.
 - Wild, F. R. W. P.; Wasiucionek, M.; Huttner, G.; Brintzinger, H. H.: Synthesis and Crystal Structure of a Chiral *ansa*-Zirconocene Derivative with Ethylene-Bridged Tetrahydroindenyl Ligands. *J. Organomet. Chem.* **288** (1985) 63-67.
 - Reddy, K. P.; Petersen, J. L.: Synthesis and Characterization of Binuclear Zirconocenophane Hydrides. The Molecular Structure of [SiMe₂(C₅H₄)₂](η^5 -C₅H₅)ZrCl(μ -H)]₂. *Organometallics* **8** (1989) 547-549.
 - Sheldrick, G. M.: Phase Annealing in SHELX-90: Direct Methods for Larger Structures. *Acta Crystallogr. A* **46** (1990) 467-473.
 - Sheldrick, G. M.: SHELXL-97. Program for the Refinement of Crystal Structures. University of Göttingen, Germany 1997.