


Crystal structure of 2,2'-[(1,4-butylene)dioxybis(nitrilomethylidyne)]-dinaphthol, C₂₆H₂₄N₂O₄

W.-K. Dong^{*,1}, J.-G. Duan¹, H.-L. Wu¹, J.-Y. Shi¹ and T.-Z. Yu^{II}

^I Lanzhou Jiaotong University, School of Chemical and Biological Engineering, Lanzhou 730070, P. R. China

^{II} Lanzhou Jiaotong University, Key Laboratory of Opto-Electronic Technology and Intelligent Control, Lanzhou 730070, P. R. China

Received November 9, 2006, accepted and available on-line December 17, 2006; CCDC no. 1267/1932



Abstract

C₂₆H₂₄N₂O₄, monoclinic, *P*12₁/*c*1 (no. 14), *a* = 5.611(2) Å, *b* = 11.723(2) Å, *c* = 16.636(3) Å, *β* = 95.432(2)°, *V* = 1089.4 Å³, *Z* = 2, *R*_{gt}(*F*) = 0.048, *wR*_{ref}(*F*²) = 0.153, *T* = 298 K.

Source of material

To an ethanol solution (10 ml) of 2-hydroxy-1-naphthaldehyde (349.5 mg, 2.01 mmol) was added an ethanol solution (3 ml) of 1,4-bis(aminooxy)butane (120.12 mg, 1.0 mmol). After the solution had been stirred at 55 °C for 4 h, the mixture was filtered, washed successively with ethanol and ethanol/hexane (1:4), respectively. The isolated compound was dried under reduced pressure and purified with recrystallization from ethanol to yield 192.8 mg of pale-brown crystalline solid (yield 45 %, m.p. 142–143 °C). Pale-brown single crystals were obtained by slow evaporation from an acetone solution at room temperature.

Elemental analysis – found: C, 72.85 %; H, 5.42 %; N, 6.73 %; calc. for C₂₆H₂₄N₂O₄: C, 72.88 %; H, 5.65 %; N, 6.54 %.

Discussion

Recently, a great deal of attention has focused on a series of *N,N'*-bis(salicylidene)ethylenediamine (salen) and its derivatives [1,2], because these compounds easily form metallosalen complexes and some of them have excellent catalytic activities for epoxidation, aziridination, etc. [3,4]. In addition, they are also used as models of reaction centers of metalloenzymes [5,6], and nonlinear optical materials [7]. Moreover, in biological and artificial systems allosteric regulation is effective in controlling molecular functions, such as molecular recognition and biological activity [8]. To tune or improve such functions, chemical modifications of a basic salen skeleton is very interesting and important [9–12]. So the introduction of some functional groups or substitution of some parts with appropriate ones are effective and inevitable. Some of such compounds have two salicylidene units, but the

linkage was limited to an alkylene (–CH=N–C–X–C–N=CH–) group due to the ease of preparation. If an *O*-alkyloxime moiety (–CH=N–O–(CH₂)_{*n*}–O–N=CH–) is used instead of (–CH=N–C–X–C–N=CH–) group, the large electronegativity of oxygen atoms is expected to affect strongly the electronic properties of N₂O₂ coordination sphere, which can lead to different and novel properties and structures of the resulted complexes. We have recently reported some *O*-alkyloxime derivatives of salen, such as 4,4'-dibromo-2,2'-[ethylenedioxybis(nitrilomethylidyne)]-diphenol [13] and the crystal structure of 4,4'-dibromo-2,2'-[(1,3-propylene)dioxybis(nitrilomethylidyne)]diphenol [14], which show high stability under the conditions where the imine derivatives suffer metathesis reaction of the C=N bonds.

The crystal structure of the title compound is built up by only the C₂₆H₂₄N₂O₄ molecules, in which all bond lengths are in normal ranges. The X-ray diffraction analysis revealed all-conformation of the (–CH=N–O–(CH₂)₄–O–N=CH–) bridge, which resulted in a planar structure with two nitrilomethylidyne units apart from each other. The IR spectrum of the compound shows a C=N stretching band at 1624 cm^{–1}, but no C=O band around 1637 cm^{–1} which is observed for 2-hydroxy-1-naphthaldehyde. The title compound is sufficiently stable to resist scrambling of the C=N bonds. This may be ascribed to lower reactivity of the oxime C=N bonds toward nucleophiles.

Table 1. Data collection and handling.

Crystal:	pale-brown block, size 0.50 × 0.53 × 0.65 mm
Wavelength:	Mo K _α radiation (0.71073 Å)
<i>μ</i> :	0.89 cm ^{–1}
Diffractionmeter, scan mode:	Bruker SMART CCD, <i>φ</i> / <i>ω</i>
2 <i>θ</i> _{max} :	50°
<i>N</i> (<i>hkl</i>) _{measured} , <i>N</i> (<i>hkl</i>) _{unique} :	5499, 1927
Criterion for <i>I</i> _{obs} , <i>N</i> (<i>hkl</i>) _{gt} :	<i>I</i> _{obs} > 2 <i>σ</i> (<i>I</i> _{obs}), 1211
<i>N</i> (<i>param</i>) _{refined} :	145
Programs:	SHELXS-97 [15], SHELXL-97 [16]

* Correspondence author (e-mail: dongwk@mail.lzjtu.cn)

Table 2. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	x	y	z	U _{iso}
H(1)	4e	1.1972	0.8157	1.5468	0.100
H(1A)	4e	1.2359	0.6307	1.7001	0.056
H(4)	4e	1.6633	0.9853	1.5794	0.069
H(5)	4e	1.9510	0.9738	1.6845	0.069
H(8)	4e	1.4947	0.6255	1.7968	0.067
H(9)	4e	1.7918	0.6162	1.8991	0.077

Table 2. Continued.

Atom	Site	x	y	z	U _{iso}
H(10)	4e	2.1167	0.7398	1.9036	0.079
H(11)	4e	2.1265	0.8795	1.8088	0.071
H(12A)	4e	0.8461	0.5936	1.4829	0.058
H(12B)	4e	0.6742	0.6705	1.5292	0.058
H(13A)	4e	0.5180	0.5057	1.5852	0.057
H(13B)	4e	0.6851	0.4303	1.5365	0.057

Table 3. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	x	y	z	U ₁₁	U ₂₂	U ₃₃	U ₁₂	U ₁₃	U ₂₃
N(1)	4e	1.0950(3)	0.6848(1)	1.5982(1)	0.049(1)	0.047(1)	0.051(1)	−0.0038(8)	0.0035(8)	−0.0048(8)
O(1)	4e	1.2990(3)	0.8659(1)	1.54699(9)	0.078(1)	0.064(1)	0.0561(9)	−0.0067(8)	−0.0068(8)	0.0108(7)
O(2)	4e	0.9297(3)	0.5962(1)	1.60133(8)	0.056(1)	0.0577(9)	0.0540(9)	−0.0124(7)	−0.0030(7)	0.0014(7)
C(1)	4e	1.2485(4)	0.6849(2)	1.6599(1)	0.051(1)	0.046(1)	0.043(1)	0.001(1)	0.006(1)	−0.0018(9)
C(2)	4e	1.4421(4)	0.7674(2)	1.6691(1)	0.047(1)	0.039(1)	0.042(1)	0.0009(9)	0.0080(9)	−0.0054(8)
C(3)	4e	1.4608(4)	0.8526(2)	1.6122(1)	0.057(1)	0.046(1)	0.043(1)	0.002(1)	0.004(1)	−0.0031(9)
C(4)	4e	1.6534(4)	0.9293(2)	1.6185(1)	0.071(2)	0.045(1)	0.057(1)	−0.005(1)	0.014(1)	0.002(1)
C(5)	4e	1.8249(4)	0.9220(2)	1.6811(1)	0.061(2)	0.048(1)	0.065(2)	−0.011(1)	0.014(1)	−0.008(1)
C(6)	4e	1.8172(4)	0.8377(2)	1.7416(1)	0.049(1)	0.047(1)	0.049(1)	0.001(1)	0.010(1)	−0.0142(9)
C(7)	4e	1.6214(4)	0.7603(2)	1.7361(1)	0.047(1)	0.041(1)	0.043(1)	0.0039(9)	0.0087(9)	−0.0087(8)
C(8)	4e	1.6207(4)	0.6772(2)	1.7980(1)	0.064(2)	0.052(1)	0.052(1)	−0.002(1)	0.002(1)	−0.000(1)
C(9)	4e	1.7990(4)	0.6711(2)	1.8591(1)	0.076(2)	0.064(2)	0.051(1)	0.006(1)	0.001(1)	0.000(1)
C(10)	4e	1.9930(5)	0.7462(2)	1.8625(1)	0.068(2)	0.075(2)	0.053(1)	0.009(1)	−0.007(1)	−0.015(1)
C(11)	4e	1.9994(4)	0.8283(2)	1.8056(1)	0.054(1)	0.065(2)	0.058(1)	−0.005(1)	0.004(1)	−0.019(1)
C(12)	4e	0.7625(3)	0.5993(2)	1.5312(1)	0.045(1)	0.050(1)	0.050(1)	0.0021(9)	0.002(1)	−0.0014(9)
C(13)	4e	0.5948(3)	0.5010(2)	1.5354(1)	0.045(1)	0.049(1)	0.049(1)	0.0042(9)	0.0043(9)	0.0001(9)

Acknowledgments. This work was supported by 'Qing Lan' Talent Engineering Funds (grant no. QL-03-01A) and Graduate Students Science Innovative Funds (grant no. DXS-2006-80) of the Lanzhou Jiaotong University, which are gratefully acknowledged.

References

- Boghaei, D. M.; Bezaatpour, A.; Behzad, M.: Synthesis, characterization and catalytic activity of novel monomeric and polymeric vanadyl Schiff base complexes. *J. Mol. Catal. A245* (2006) 12–16.
- Abu-Surrah, A. S.; Thewalt, U.; Rieger, B.: Chiral palladium(II) complexes bearing tetradentate nitrogen ligands: synthesis, crystal structure and reactivity towards the polymerization of norbornene. *J. Organomet. Chem.* **587** (1999) 58–66.
- Jacobsen, E. N.; Zhang, W.; Muci, A. R.; Ecker, J. R.; Deng, L.: Highly enantioselective epoxidation catalysts derived from 1,2-diaminocyclohexane. *J. Am. Chem. Soc.* **113** (1991) 7063–7064.
- Venkataramanan, N. S.; Kuppuraj, G.; Rajagopal, S.: Metal-salen complexes as efficient catalysts for the oxygenation of heteroatom containing organic compounds-synthetic and mechanistic aspects. *Coord. Chem. Rev.* **249** (2005) 1249–1268.
- Tsou, T. T.; Loots, M.; Halpern, J.: Kinetic determination of transition metal-alkyl bond dissociation energies: application to organocobalt compounds related to B₁₂ coenzymes. *J. Am. Chem. Soc.* **104** (1982) 623–624.
- Summers, M. F.; Marzilli, L. G.; Bresciani-Pahor, N.; Randaccio, L.: Unusual bond lengths, conformations, and ligand exchange rates in B12 models with the bis(salicylidene)-o-phenylenediamine equatorial ligand. *J. Am. Chem. Soc.* **106** (1984) 4478–4485.
- Lacroix, P. G.: Second-order optical nonlinearities in coordination chemistry: The case of bis(salicylaldiminato)metal Schiff base complexes. *Eur. J. Inorg. Chem.* (2001) 339–348.
- Sun, S. S.; Stern, C. L.; Nguyen, S. T.; Hupp, J. T.: Directed assembly of transition-metal-coordinated molecular loops and square from salen-type components. Examples of metalation-controlled structural conversion. *J. Am. Chem. Soc.* **126** (2004) 6314–6326.
- Akine, S.; Taniguchi, T.; Nabeshima, T.: Cooperative formation of trinuclear zinc(II) complexes via complexation of a tetradentate oxime chelate ligand, Salamo, and zinc(II) acetate. *Inorg. Chem.* **43** (2004) 6142–6144.
- Akine, S.; Takanori, T.; Dong, W. K.; Nabeshima, T.: Oxime-Based Salen-Type Tetradentate Ligands with High Stability against Imine Metathesis Reaction. *J. Org. Chem.* **70** (2005) 1704–1711.
- Akine, S.; Dong, W. K.; Nabeshima, T.: Octanuclear zinc(II) and cobalt(II) clusters produced by cooperative tetrameric assembling of oxime chelate ligands. *Inorg. Chem.* **45** (2005) 4677–4684.
- Akine, S.; Taniguchi, T.; Nabeshima, T.: Novel synthetic approach to trinuclear 3d – 4f complexes: specific exchange of the central metal of a trinuclear zinc(II) complex of a tetraoxime ligand with a lanthanide(III). *Angew. Chem., Int. Ed.* **41** (2002) 4670–4673.
- Dong, W. K.; Feng, J. H.: 4,4'-Dibromo-2,2'-[ethylenedioxybis(nitrilomethylidyne)]diphenol. *Acta Crystallogr. E62* (2006) o3577–o3578.
- Dong, W. K.; Feng, J. H.; Yang, X.-Q.: Crystal structure of 4,4'-dibromo-2,2'-[(1,3-propylene)dioxybis(nitrilomethylidyne)]diphenol, C₁₇H₁₆Br₂N₂O₄. *Z. Kristallogr. NCS* **221** (2006) 447–448.
- Sheldrick, G. M.: SHELXS-97. Program for the Solution of Crystal Structures. University of Göttingen, Germany 1997.
- Sheldrick, G. M.: SHELXL-97. Program for the Refinement of Crystal Structures. University of Göttingen, Germany 1997.