

Zeitschrift für Kristallographie

New Crystal Structures

Volume 221 · Issue 1 · 2006

Imorganic Crystal Structures

V. Babizhetkyy, Hj. Mattausch and A. Simon

Crystal structure of pentapraseodymium diboropentacarbide,
 $\text{Pr}_{55}\text{B}_2\text{C}_5$ 1

M. Lukachuk, Hj. Mattausch and A. Simon

Crystal structure of tetragadolinium hexaiodide
monogermanide, $\text{Gd}_4\text{I}_6\text{Ge}$ 3

Organic and Metalorganic Crystal Structures

X. Li

Crystal structure of bis(1,10-phenanthroline-*N,N'*)bis(4-fluorobenzoato-*O,O'*)tetrakis(μ -4-fluorobenzoato-*O,O'*)-diineodymium(III), $\text{Nd}_2(\text{C}_6\text{H}_4\text{FCOO})_6(\text{C}_{12}\text{H}_8\text{N}_2)_2$ 5

Y.-L. Ju, X. Li and T.-T. Zhang

Crystal structure of bis(1,10-phenanthroline-*N,N'*)bis(4-fluorobenzoato-*O,O'*)tetrakis(μ -4-fluorobenzoato-*O,O'*)-di dysprosium(III), $\text{Dy}_2(\text{C}_6\text{H}_4\text{FCOO})_6(\text{C}_{12}\text{H}_8\text{N}_2)_2$ 7

T.-T. Zhang, X. Li and Y.-L. Ju

Crystal structure of bis(1,10-phenanthroline-*N,N'*)bis(2,4-dimethylbenzoato-*O,O'*)tetrakis(μ -2,4-dimethylbenzoato-*O,O'*)disamarium(III), $\text{Sm}_2(\text{C}_9\text{H}_9\text{O}_2)_6(\text{C}_{12}\text{H}_8\text{N}_2)_2$ 9

A. Al-Harrasi, I. Brüdgam, H. Hartl and H.-U. Reißig

Crystal structure of $3S,4'S,6'S\text{-}2\text{-benzyl-3-[}(tert\text{-butyl-dimethyl-siloxy)}\text{-}(2',2'\text{-dimethyl-1,3-dioxolan-4'-yl)}\text{-}6'\text{-methyl]}4\text{-[}2\text{-[(trimethylsilyl)ethoxy]}3\text{-6-dihydro-2H-1,2-oxazine, C}_{28}\text{H}_{49}\text{NO}_5\text{Si}_2$ 12

M. Zeller, S. J. DiMuzio, K. Bee, J. Clemente, J. Frederick,

A. Fucci, L. Gordon, A. Harper, A. Kasmar, C. Kasmar,

S. Polomsky, A. Simmons, D. Smallsreed, T. Underwood

and A. D. Hunter

Crystal structures of the methylene chloride solvates of tetrakis(3,5-dimethoxyphenyl)porphyrin and its isotopic copper(II) and nickel(II) complexes, $\text{C}_{52}\text{H}_{46}\text{N}_4\text{O}_8 \cdot 2\text{CH}_2\text{Cl}_2$, $\text{Cu}(\text{C}_{52}\text{H}_{44}\text{N}_4\text{O}_8) \cdot 2\text{CH}_2\text{Cl}_2$, $\text{Ni}(\text{C}_{52}\text{H}_{44}\text{N}_4\text{O}_8) \cdot 2\text{CH}_2\text{Cl}_2$ 15

T. Irrgang, S. Karthikeyan, S. Balireddi and R. Kempe

Crystal structure of *N,N'*-bis(2-methyl-4,5-diphenyl-imidazo[1,5-*b*]pyridazin-7-yl)-1,3-diaminopropane ethanol solvate, $\text{C}_{32}\text{H}_{6}(\text{NH})_2(\text{C}_{19}\text{H}_{14}\text{N}_3)_2 \cdot \text{C}_2\text{H}_5\text{OH}$ 20

T. Irrgang, S. Balireddi, S. Karthikeyan and R. Kempe

Crystal structure of *N,N'*-bis(2-methyl-4-phenyl-5-*tert*-butylimidazo[1,5-*b*]pyridazin-7-yl)ethylenediamine, $\text{C}_{22}\text{H}_{4}(\text{NH})_2(\text{C}_{17}\text{H}_{18}\text{N}_3)_2$ 23

D.-P. He, Y.-L. Dou and J.-B. She

Crystal structure of 1,6-bis(1-phenyl-1*H*-tetrazol-5-ylthio)hexane, $\text{C}_6\text{H}_{12}(\text{SCN}_4\text{C}_6\text{H}_5)_2$ 25

Y.-L. Wang, W.-T. Yu, D. Xu, X.-Q. Wang, W.-F. Guo and G.-H. Zhang

Crystal structure of bis(*N*-ethylpyridinium) bis(2-thioxo-1,3-dithiole-4,5-dithiolato)mercurate(II), $(\text{C}_7\text{H}_{10}\text{N})_2[\text{Hg}(\text{C}_3\text{S}_5)_2]$ 27

L.-F. Zhao

Crystal structure of (*E*)-4-methoxy-nitrophenone *N*-(2,4-dinitrophenyl)hydrazone, $\text{C}_{14}\text{H}_{11}\text{N}_5\text{O}_7$ 29

P. Kuhn, D. Matt, L. Toupet and A. Louati

Crystal structure of 25,27-dibenzyloxy-26-benzoyloxy-28-hydroxycalix[4]arene, $\text{C}_{49}\text{H}_{40}\text{O}_5$ 31

L. Wiehl, J. Schreuer and E. Haussührl

Crystal structure of poly-[*[(3-hydroxypyridine)oxalato-copper(II)] monohydrate*, $\text{Cu}(\text{NC}_5\text{H}_4\text{OH})(\text{COO})_2 \cdot \text{H}_2\text{O}$ 35

J. Xiong, X.-X. Lei, M.-L. Hu, J.-X. Yuan and X.-Q. Cai

Crystal structure of 2-(5-fluoro-2,4-dioxo-1,2,3,4-tetrahydropyrimidine)glycin, $\text{C}_8\text{H}_8\text{FN}_3\text{O}_5$ 37

M.-L. Hu, P. Yin, J. Xiong and J.-X. Yuan

Crystal structure of 2(*R*)-(5-fluorouracil-1-yl)acetyl-2-phenylglycin dimethylformamide solvate, $\text{C}_{14}\text{H}_{12}\text{FN}_3\text{O}_5 \cdot \text{C}_3\text{H}_7\text{NO}$ 39

M.-F. Li and Y.-K. Shan

Crystal structure of *trans*-bis(triphenylphosphine oxide)-tetrachlorotitanium(IV), $\text{TiCl}_4[(\text{C}_6\text{H}_5)_3\text{PO}]_2$ 41

W. Frey, E. Haug and W. Kantlehner

Crystal structure of 1-(4-chlorophenyl)-2-(5-methyl-1,3,4-thiadiazol-2-yl)ethanone, $\text{C}_{11}\text{H}_9\text{ClN}_2\text{OS}$ 43

A. Bekaert, P. Lemoine, J. D. Brion and B. Viossat

Crystal structure of bis(acetato-*O,O'*)bis(μ -acetato-*O,O'*)bis(κ^3 *N*-antipyrine-*O*)tetraoxodiuranium, $\text{U}_2\text{O}_4(\text{CH}_3\text{COO})_4[\text{C}_6\text{H}_5\text{N}_2\text{C}_3(\text{CH}_3)_2\text{HO}]_2$ 45

M.-L. Hu and F. Chen

Crystal structure of bis(2,2'-biimidazol-1-iun) tetrachloroferrate(III) chloride, $(\text{C}_6\text{H}_7\text{N}_4)_2[\text{FeCl}_4]\text{Cl}$ 47

T. L. Balliano, M. A. Pereira, C. A. De Simone, V. R. S. Malta,

N. Velasquez, A. G. Cioletti and M. O. F. Goulart

Crystal structure of 1-(2-bromoethyl)-2-methyl-5-nitroimidazole, $\text{C}_6\text{H}_8\text{BrN}_3\text{O}_2$ 49

| | |
|---|-----|
| <i>C. Wang, Y.-Q. Hou, J. Li and F.-X. Zhang</i> Crystal structure of methanol-[<i>N,N'</i> -o-phenylenebis(salicyl- aldiminato- <i>N,N',O,O'</i>)(salicylaldehydato-O)manganese(III), $\text{Mn}(\text{CH}_3\text{OH})(\text{C}_6\text{H}_4\text{OCHO})[\text{C}_6\text{H}_4(\text{C}_6\text{H}_4\text{OCHN})_2]_2$ | 51 |
| <i>J.-Y. Ma, T.-X. Wu, X.-G. She and X.-F. Pan</i> Crystal structure of dichloro((<i>N</i> -(<i>N',N'</i> -dimethylammonio)- ethyl)-2,4-dichlorosalicylaliminato- <i>N,O</i>)copper(II), $\text{Cu}(\text{C}_{11}\text{H}_{14}\text{Cl}_2\text{N}_2\text{O})\text{Cl}_2$ | 53 |
| <i>T.-B. Li, D. Xu, W.-T. Yu, G.-H. Zhang and X.-Q. Wang</i> Crystal structure of hexadecyltrimethylammonium bis(1,3-dithiole-2-thione-4,5-dithiolato)aurate(III), $[\text{C}_{16}\text{H}_{33}(\text{CH}_3)_3\text{N}][\text{Au}(\text{C}_3\text{S}_5)_2]$ | 55 |
| <i>L.-J. Zhang, J.-C. Ding, H.-Y. Wu and M.-C. Liu</i> Crystal structure of 5-fluoro-1-(carboxymethyl)uracil, $\text{C}_6\text{H}_5\text{FN}_2\text{O}_4$ | 57 |
| <i>L.-J. Zhang, M.-C. Liu, J.-C. Ding and H.-Y. Wu</i> Crystal structure of theophylline-7-acetic acid, $\text{C}_9\text{H}_{10}\text{N}_4\text{O}_4$ | 59 |
| <i>M.-G. Zhao</i> Crystal structure of tetraaquabis(4-nitropyridine <i>N</i> -oxide- <i>O</i>)- nickel(II) dinitrate, $[\text{Ni}(\text{H}_2\text{O})_4(\text{ONC}_5\text{H}_4\text{NO}_2)_2][\text{NO}_3]_2$ | 61 |
| <i>Y.-Y. Yu</i> Crystal structure of <i>N,N'</i> -bis(salicylidene)-1,10- decanediamine, $\text{C}_{10}\text{H}_{20}(\text{NCHC}_6\text{H}_4\text{OH})_2$ | 63 |
| <i>Z.-C. Ma</i> Crystal structure of (<i>R</i>)-1,1'-bi-2-naphthol 1,3-bis(pyrid-4-yl)propane, $\text{C}_{20}\text{H}_{14}\text{O}_2 \cdot \text{C}_{13}\text{H}_{14}\text{N}_2$ | 65 |
| <i>B. Tinant, M. Laurent and J. Marchand-Brynaert</i> Crystal structure of <i>N</i> -(diphenyl)methyl- <i>N</i> -(3',3'-dimethyl- 2'-oxo)butyl-2(<i>R</i>),3(<i>R</i>)-epoxybutyramide, $\text{C}_{23}\text{H}_{27}\text{NO}_3$ | 68 |
| <i>B. Tinant, M. Laurent and J. Marchand-Brynaert</i> Crystal structure of <i>N</i> -(diphenyl)methyl-3(<i>S</i>)-[1'(<i>R</i>)- hydroxoethyl]-4(<i>S</i>)-[2'',2''-dimethyl-1''-oxopropyl]- 2-azetidinone, $\text{C}_{23}\text{H}_{27}\text{NO}_3$ | 71 |
| <i>X.-Q. Han and G.-R. Yue</i> Crystal structure of bis[N-cyclopropyl-5-nitrosalicyl- aldiminato]copper(II), $\text{Cu}[(\text{C}_3\text{H}_5\text{NCH})(\text{O})\text{C}_6\text{H}_3(\text{NO}_2)]_2$ | 73 |
| <i>J. Kaizer, S. Góger, G. Speier, M. Réglier and M. Giorgi</i> Crystal structure of μ -dichlorobis(<i>N</i> -benzyl-bis(2-pyridyl- methyl)amino)dicopper(II) diperchlorate ethanol solvate (1:2%), $[\{\text{Cu}_2\text{Cl}_2\{\text{N}(\text{C}_6\text{H}_6\text{N})_2(\text{C}_7\text{H}_7)\}_2\}_2\text{ClO}_4]_2 \cdot \frac{1}{2}\text{C}_2\text{H}_5\text{OH}$ | 75 |
| <i>L. Wiehl, J. Schreuer and E. Haussühl</i> Crystal structures of catena-(tris(μ -betaine- <i>O,O'</i>)- manganese(II)) tetrachlorozincate and catena- (tris(μ -betaine- <i>O,O'</i>)manganese(II)) tetrachlorocobaltate, $[(\{\text{CH}_3\}_3\text{NCH}_2\text{COO}\}_3\text{Mn}][\text{MCl}_4]$ ($\text{M} = \text{Zn}, \text{Co}$) | 77 |
| <i>X.-Q. Zhang, Z.-T. Zhang and X.-L. Zhang</i> Crystal structure of kakkalidone, $\text{C}_{23}\text{H}_{24}\text{O}_{11}$ | 80 |
| <i>J.-C. Ding, M.-C. Liu, L.-J. Zhang and H.-Y. Wu</i> Crystal structure of 5-fluoro-1-benzyluracil, $(\text{C}_6\text{H}_5)\text{CH}_2(\text{C}_4\text{N}_2\text{H}_2\text{FO}_2)$ | 83 |
| <i>L. Tang, J. Li, D.-S. Li, C.-H. Zhou, J.-J. Wang and J.-W. Wang</i> Crystal structure of bis(2-hydroxy-2,2-bis(pyridin- 2-yl)acetato)nickel(II) methanol solvate, $\text{Ni}[(\text{C}_5\text{H}_4\text{N})_2\text{C}(\text{OH})\text{COO}]_2 \cdot \text{CH}_3\text{OH}$ | 85 |
| <i>W. Frey, D. Shaw, A. Jégou and V. Jäger</i> Crystal structure of dimethyl (2 <i>S</i> ,3 <i>R</i> ,3 <i>aS</i> ,4 <i>S</i> ,5 <i>R</i> ,6 <i>R</i>)- 6-iodomethyl-4,5-isopropylidenedioxy-hexahydro- pyrrolo[1,2- <i>b</i>]pyrazole-2,3-dicarboxylate, $\text{C}_{14}\text{H}_{21}\text{IN}_2\text{O}_6$ | 87 |
| <i>W. Frey, M. Gulla and V. Jäger</i> Crystal structure of <i>rel</i> -(2 <i>R</i> ,3 <i>aS</i>)-2-bromomethyl- 3,3 <i>a</i> ,4,5,6,7-hexahydro-2 <i>H</i> -indole 1-oxide, $\text{C}_9\text{H}_{14}\text{BrNO}$ | 89 |
| <i>S. E. McKay, K. A. Wheeler and B. Holthouse</i> Crystal structure of 2,2':6',2"-terpyridine 1,1"-dioxide 1,2,4,5-tetracyanobenzene, $\text{C}_{15}\text{H}_{11}\text{N}_3\text{O}_2 \cdot \text{C}_6\text{H}_4(\text{CN})_4$ | 91 |
| <i>S. Deeken, T. Irrgang and R. Kempe</i> Crystal structure of 1,5-cyclooctadiene-bis[(4-methyl- pyridin-2-yl)-trimethylsilylanyl-amido]ruthenium(II), $\text{Ru}(\text{C}_8\text{H}_{12})(\text{C}_9\text{H}_{15}\text{N}_2\text{Si})_2$ | 93 |
| <i>T. Schmalz, S. Karthikeyan, T. Irrgang and R. Kempe</i> Crystal structure of <i>cis</i> -bis[2-(diphenylphosphinoaminato)- 4-methylpyridine]platinum(II), $\text{Pt}(\text{C}_{18}\text{H}_{16}\text{N}_2\text{P})_2$ | 95 |
| <i>M. Buchta, A. Jegorov, M. Hušák, J. Cejka, B. Kratochvíl, M. Kuzma and P. Sedmera</i> Crystal structure of taxuspinanane A acetonitrile solvate, $\text{C}_{47}\text{H}_{59}\text{NO}_{14} \cdot \text{CH}_3\text{CN}$ | 97 |
| <i>Q.-W. Zhang and G.-X. Wang</i> Crystal structure of catena-[dodecaaqua(μ^3 -benzene- 1,3,5-tricarboxylato)(μ^2 -benzene-1,3,5-tricarboxylato)- tricopper(II)], $\text{Cu}_3(\text{H}_2\text{O})_{12}(\text{C}_9\text{H}_3\text{O}_6)_2$ | 101 |
| <i>G. Laus, C. E. Strasser, K. Wurst and H. Schottenberger</i> Crystal structure of (<i>E</i>)-2-ruthenocenylethenylcobaltocenium hexafluorophosphate, $[(\text{C}_5\text{H}_5)\text{Ru}(\text{C}_{12}\text{H}_{10})\text{Co}(\text{C}_5\text{H}_5)][\text{PF}_6]$ | 103 |
| <i>Y. He, X.-L. Cheng and Y. Chang</i> Crystal structure of methylammonium 4',7-dimethoxy- isoflavone-3'-sulfonate monohydrate, $(\text{CH}_3\text{NH}_3)(\text{C}_{17}\text{H}_{13}\text{O}_4\text{SO}_3) \cdot \text{H}_2\text{O}$ | 105 |

Instructions to Contributors

Author Index

Formulae Index

Further details of the structure determination can be accessed at the relevant database:

Data with CSD numbers can be obtained from the Fachinformationszentrum Karlsruhe, Gesellschaft für wissenschaftlich-technische Information mbH, D-76344 Eggenstein-Leopoldshafen, Germany (e-mail: docdel@fiz-karlsruhe.de).

Data with CCDC numbers can be obtained from the Cambridge Crystallographic Data Centre (CCDC), 12 Union Road, Cambridge CB2 1EZ, UK, (fax: +44-(0)1223-33 60 33 or e-mail: deposit@chemcrys.cam.ac.uk).