

Crystal structure of (1*S*,2*R*,4*S*)-1-((phenylamino)methyl)-4-(prop-1-en-2-yl)cyclohexane-1,2-diol, C₁₆H₂₃NO₂


Rachid Outouch^I, Brahim Boualy^I, Larbi El Firdoussi^{*I}, Mustapha Ait Ali^I, Corrado Rizzoli^{II}
and Anke Spannenberg^{III}

¹ Laboratoire de Chimie de Coordination, Faculté des Sciences-Semlalia BP 2390, 40001 Marrakech, Morocco

^{II} Dipartimento di Chimica Generale ed Inorganica, Chimica Analitica, Chimica Fisica, Università degli Studi di Parma, 43100 Parma, Italy

^{III} Leibniz-Institut für Katalyse e. V. an der Universität Rostock, Albert-Einstein-Straße 29a, 18059 Rostock, Germany

Received February 9, 2011, accepted and available on-line April 27, 2011; CCDC no. 1267/3399



Abstract

$\text{C}_{16}\text{H}_{23}\text{NO}_2$, orthorhombic, $P2_12_12_1$ (no. 19), $a = 5.9637(3)$ Å, $b = 8.8317(5)$ Å, $c = 27.809(1)$ Å, $V = 1464.7$ Å³, $Z = 4$, $R_{\text{ref}}(F) = 0.026$, $wR_{\text{ref}}(F^2) = 0.040$, $T = 150$ K.

Source of material

A mixture of aniline (5.1 mmol) and epoxy perillyl alcohol (5 mmol) was added to 0.25 mmol of $\text{Ca}(\text{CF}_3\text{CO}_2)_2$ under solvent-free conditions. The mixture was stirred at 40 °C for 72 h. After the reaction was finished, the mixture was extracted with AcOEt (3 × 10 mL), dried over Na_2SO_4 and the solvent was removed at reduced pressure. Colorless crystals of the title compound were isolated by column chromatography on silica gel using hexane/ethyl acetate as eluent (yield 55 %).

Experimental details

Hydrogen atoms attached to N1, O1 and O2 were located in the difference Fourier map and refined isotropically. All other hydrogen atoms were placed in idealized positions and refined using a riding model. The absolute configuration of the title structure could not be determined using data collected with Mo K_{α} radiation (Flack parameter $x = 0.1(10)$).

Discussion

Aminodiols are present in many pharmacologically important compounds such as antiviral glycosidase inhibitors [1,2], sphingolipids [3,4], antibacterial and antiprotozoal [5]. Also, aminodiol has been found to act as HIV protease inhibitors [6,7]. Furthermore, aminodiol may serve as useful starting materials for the synthesis of biologically active compounds, e.g., 3-amino-1,2-butanediol derivative, a key intermediate in the synthesis of anticancer agent ES-285 [8]. Chiral aminodiol and their derivatives also find application as catalysts for enantioselective transformations [9,10]. Different approaches to obtain aminodiols have been developed [11-13]. Aminolysis of 1,2-

epoxides represents one of the valuable pathway to produce commercially important aminoalcohols and aminodiols from olefins [14-16].

The absolute configuration has been assigned by reference to unchanging the chiral centres in the synthetic procedure. The cyclohexane ring is in a chair conformation, with puckering parameters Q , θ and φ of 0.565(2) Å, 172.8(1)° and -31(1)°, respectively [19]. In the crystal structure, molecules interact *via* N1–H1A...O2 and O2–H2...O1 intermolecular hydrogen bonds to form chains parallel to [100]. The chains are further linked into layers parallel to the (001) plane by O1–H1B...N1 hydrogen bonds.

Table 1. Data collection and handling.

Crystal:	colorless needle, size 0.08 × 0.16 × 0.47 mm
Wavelength:	Mo K α radiation (0.71073 Å)
μ :	0.77 cm ⁻¹
Diffractometer, scan mode:	STOE IPDS II, ω
2 θ _{max} :	51°
$N(hkl)$ _{measured} , $N(hkl)$ _{unique} :	9453, 2721
Criterion for I_{obs} , $N(hkl)$ _{gt} :	$I_{\text{obs}} > 2 \sigma(I_{\text{obs}})$, 1941
$N(\text{param})$ _{refined} :	185
Programs:	SHELXS-97, SHELXL-97 [20]

Table 2. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	x	y	z	U_{iso}
H(1C)	4a	0.2794	0.5111	0.7883	0.031
H(1D)	4a	0.2056	0.3386	0.7803	0.031
H(3)	4a	0.1020	0.6135	0.7143	0.031
H(4A)	4a	0.2151	0.5950	0.6350	0.034
H(4B)	4a	-0.0472	0.5571	0.6367	0.034
H(5)	4a	0.0329	0.2945	0.6332	0.035
H(6A)	4a	0.4090	0.2120	0.6216	0.034
H(6B)	4a	0.4948	0.3831	0.6273	0.034
H(7A)	4a	0.2723	0.2133	0.7013	0.031
H(7B)	4a	0.5337	0.2543	0.7015	0.031
H(9A)	4a	-0.0613	0.3247	0.5112	0.060
H(9B)	4a	-0.1539	0.2582	0.5628	0.060
H(10A)	4a	0.2597	0.4718	0.5064	0.078

* Correspondence author (e-mail: elfirdoussi@ucam.ac.ma)

Table 2. Continued.

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso}
H(10B)	4a	0.4439	0.4259	0.5456	0.078
H(10C)	4a	0.2964	0.5755	0.5529	0.078
H(12)	4a	0.2648	0.3128	0.8642	0.039
H(13)	4a	0.3373	0.3287	0.9463	0.049
H(14)	4a	0.6854	0.4077	0.9739	0.054

Table 2. Continued.

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso}
H(15)	4a	0.9675	0.4677	0.9189	0.046
H(16)	4a	0.8995	0.4484	0.8368	0.037
H(1A)	4a	0.653(2)	0.408(2)	0.7756(5)	0.033(4)
H(1B)	4a	0.480(3)	0.642(2)	0.7119(6)	0.066(5)
H(2)	4a	−0.187(3)	0.457(2)	0.7094(6)	0.068(7)

Table 3. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> ₁₁	<i>U</i> ₂₂	<i>U</i> ₃₃	<i>U</i> ₁₂	<i>U</i> ₁₃	<i>U</i> ₂₃
C(1)	4a	0.3223(2)	0.4147(2)	0.77270(4)	0.0214(8)	0.0232(7)	0.0318(8)	0.0017(7)	−0.0006(6)	−0.0007(6)
C(2)	4a	0.3326(2)	0.4377(1)	0.71836(5)	0.0195(7)	0.0199(7)	0.0299(7)	−0.0030(7)	0.0024(7)	0.0027(6)
C(3)	4a	0.1144(2)	0.5092(2)	0.70053(4)	0.0206(8)	0.0213(7)	0.0350(8)	−0.0010(7)	0.0014(7)	0.0012(6)
C(4)	4a	0.1032(2)	0.5201(1)	0.64617(4)	0.0248(9)	0.0263(8)	0.0327(8)	0.0006(7)	−0.0050(7)	0.0018(6)
C(5)	4a	0.1479(3)	0.3681(1)	0.62143(5)	0.031(1)	0.0258(8)	0.0320(8)	−0.0021(8)	−0.0043(7)	0.0010(6)
C(6)	4a	0.3775(3)	0.3104(2)	0.63731(4)	0.029(1)	0.0219(7)	0.0326(8)	0.0024(7)	−0.0016(7)	−0.0027(6)
C(7)	4a	0.3834(3)	0.2910(1)	0.69181(4)	0.0237(9)	0.0217(7)	0.0327(8)	0.0005(7)	−0.0033(6)	−0.0005(6)
C(8)	4a	0.1223(3)	0.3804(2)	0.56749(5)	0.050(1)	0.0254(8)	0.0343(9)	0.0054(8)	−0.0099(8)	−0.0021(6)
C(9)	4a	−0.0467(3)	0.3150(2)	0.54505(6)	0.064(1)	0.046(1)	0.0415(9)	0.006(1)	−0.0184(9)	−0.0046(8)
C(10)	4a	0.2957(3)	0.4714(2)	0.54075(6)	0.078(2)	0.045(1)	0.0331(8)	−0.002(1)	−0.0023(9)	0.0054(8)
C(11)	4a	0.5742(3)	0.3803(1)	0.84209(5)	0.0306(9)	0.0199(7)	0.0263(8)	0.0047(7)	−0.0023(7)	−0.0003(5)
C(12)	4a	0.4084(3)	0.3443(2)	0.87508(5)	0.030(1)	0.0356(9)	0.0310(8)	0.0009(8)	−0.0005(7)	0.0003(6)
C(13)	4a	0.4516(3)	0.3542(2)	0.92388(5)	0.043(1)	0.048(1)	0.0319(9)	0.0099(9)	0.0038(8)	0.0020(7)
C(14)	4a	0.6578(3)	0.4003(2)	0.94035(5)	0.056(1)	0.049(1)	0.0301(8)	0.018(1)	−0.0091(9)	−0.0082(7)
C(15)	4a	0.8246(3)	0.4358(2)	0.90779(5)	0.039(1)	0.0336(9)	0.044(1)	0.0064(9)	−0.0157(8)	−0.0079(7)
C(16)	4a	0.7836(3)	0.4250(2)	0.85906(5)	0.0286(9)	0.0264(8)	0.0382(9)	0.0027(8)	−0.0025(7)	−0.0033(7)
O(1)	4a	0.5139(2)	0.5399(1)	0.70720(3)	0.0204(6)	0.0203(5)	0.0360(5)	−0.0016(4)	0.0012(4)	0.0012(4)
O(2)	4a	−0.0653(2)	0.4193(1)	0.71907(4)	0.0208(6)	0.0378(6)	0.0437(6)	−0.0029(5)	−0.0002(5)	0.0081(5)
N(1)	4a	0.5391(2)	0.3634(1)	0.79207(4)	0.0212(7)	0.0278(6)	0.0265(7)	0.0000(6)	0.0012(6)	0.0027(5)

References

- Hughes, A. B.; Rudge, A. J.: Deoxynojirimycin: synthesis and biological activity. *Nat. Prod. Rep.* **11** (1994) 135–162.
- Liang, P. H.; Cheng, W. C.; Lee, Y. L.; Yu, H. P.; Wu, Y. T.; Lin, Y. L.; Wong, C. H.: Novel five-membered iminocyclitol derivatives as selective and potent glycosidase inhibitors: new structures for antivirals and osteoarthritis. *ChemBioChem.* **7** (2006) 165–173.
- Hannun, Y. A.; Obeid, L. M.: The Ceramide-centric universe of lipid-mediated cell regulation: stress encounters of the lipid kind. *J. Biol. Chem.* **277** (2002) 25847–25850.
- Nakamura, T.; Shiozaki, M.: Stereoselective synthesis of D-erythro-sphingosine and L-lyxo-phytosphingosine. *Tetrahedron* **57** (2001) 9087–9092.
- Gondela, E.; Walczak, K. Z.: Synthesis and preliminary bioactivity assays of 3,4-dichloro-5-(*ω*-hydroxyalkylamino)-2(5*H*)-furanones. *Eur. J. Med. Chem.* **45** (2010) 3993–3997.
- Kempf, D. J.; Sowin, T. J.; Doherty, E. M.; Hannick, S. M.; Codavoci, L.; Henry, R. F.; Green, B. E.; Spanton, S. G.; Norbeck, D. W.: Stereocontrolled synthesis of C₂-symmetric and pseudo-C₂-symmetric diamino alcohols and diols for use in HIV protease inhibitors. *J. Org. Chem.* **57** (1992) 5692–5700.
- Wang, G. T.; Li, S.; Wideburg, N.; Krafft, G. A.; Kempf, D. J.: Synthetic chemical diversity: solid phase synthesis of libraries of C₂ symmetric inhibitors of HIV protease containing diamino diol and diamino alcohol cores. *J. Med. Chem.* **38** (1995) 2995–3002.
- Allepuz, A. C.; Badorrey, R.; Daz-de-Villegas, M. D.; Gálvez, J. A.: Diastereoselective reduction of ketimines derived from (*R*)-3,4-dihydroxybutan-2-one: an alternative route to key intermediates for the synthesis of anticancer agent ES-285. *Tetrahedron: Asymmetry* **21** (2010) 503–506.
- Panev, S.; Linden, A.; Dimitrov, V.: Chiral aminoalcohols with a menthane skeleton as catalysts for the enantioselective addition of diethylzinc to benzaldehyde. *Tetrahedron: Asymmetry* **12** (2001) 1313–1321.
- Braga, A. L.; Rubim, R. M.; Schrekker, H. S.; Wessjohann, L. A.; de Bolster, M. W. G.; Zeni, G.; Sehnem, J. A.: The facile synthesis of chiral oxazoline catalysts for the diethylzinc addition to aldehydes. *Tetrahedron: Asymmetry* **14** (2003) 3291–3295.
- Kwon, S. J.; Ko, S. Y.: Stereoselective Synthesis of triply-stereogenic syn, anti-amino diols: the Abbott amino diol. *Bull. Korean Chem. Soc.* **24** (2003) 1053–1054.
- Dias, L. C.; Fattori, J.; Perez, C. C.; de Oliveira, V. M.; Aguilar, A. M.: An efficient procedure for the synthesis of 2-N-Boc-amino-3,5-diols. *Tetrahedron* **64** (2008) 5891–5903.
- Szakonyi, Z.; Hetenyi, A.; Fulop, F.: Synthesis and application of monoterpene-based chiral aminodiols. *Tetrahedron* **64** (2008) 1034–1039.
- Fan, R. H.; Hou, X. L.: Efficient ring-opening reaction of epoxides and aziridines promoted by tributylphosphine in water. *J. Org. Chem.* **68** (2003) 726–730.
- Kamal, A.; Ramu, R.; Amerudin, M.; Azhar, G. B.; Khanna, R.: Copper(II) tetrafluoroborate-catalyzed ring-opening of epoxides by amines. *Tetrahedron Lett.* **46** (2005) 2675–2677.
- Carree, F.; Gil, R.; Collin, J.: Samarium iodides catalyzed meso-epoxides ring opening by aromatic amines. *Tetrahedron Lett.* **45** (2004) 7749–7751.
- Harrad, M. A.; Outtouch, R.; Ait Ali, M.; El Firdoussi, L.; Karim, A.; Roucoux, A.: Ca(CF₃CO₂)₂: An efficient Lewis acid catalyst for chemo- and regio-selective enamination of β-dicarbonyl compounds. *Catal. Commun.* **11** (2010) 442–446.
- Bach, R. D.; Klein, M. W.; Ryntz, R. A.; Holubka, J. W.: Epoxidation of alkenes with O-ethylperoxycarbonic acid generated *in situ* in an alkalineiphase solvent system. *J. Org. Chem.* **14** (1979) 2569–2571.
- Cremer, D.; Pople, J. A.: General definition of ring puckering coordinates. *J. Am. Chem. Soc.* **97** (1975) 1354–1358.
- Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr.* **A64** (2008) 112–122.