

Spontaneous contralateral pneumothorax in a patient with low Body Mass Index

Case Report

Kensuke Nakazawa, Gen Ohara, Katsunori Kagohashi,
Koichi Kurishima, Atsushi Ishibashi, Hiroaki Satoh

*Division of Respiratory Medicine, Mito Medical Center, University of Tsukuba
Miya-machi 3-2-7, Mito, Ibaraki 310-0015, Japan*

Received 12 March 2012; Accepted 28 August 2012

Abstract: Spontaneous pneumothorax is most common in adolescents and young adults. Some of them develop contralateral pneumothorax. In this paper, we report the case of a patient with spontaneous contralateral pneumothorax, whose body mass index (BMI) was 18.8 kg/m². For either chest physicians or thoracic surgeons, follow up with recognition of increased risk of the contralateral pneumothorax is important especially in patients with contralateral bullous lesions and low BMI.

Keywords: Spontaneous pneumothorax • Contralateral pneumothorax • Body mass index • Video-assisted thoracic surgery

© Versita Sp. z o.o.

1. Introduction

Spontaneous pneumothorax is troublesome for patients, due to its tendency to recur. The average rate of ipsilateral recurrence of spontaneous pneumothorax is 30% (range, 16 to 52%) [1], and the contralateral occurrence rate is around 15% [2-6]. Herein, we present a case of contralateral pneumothorax, which was successfully treated by video-assisted thoracoscopic surgery (VATS).

2. Case report

An 18-year-old man presented to our hospital with sudden onset of right chest pain. He had been in good health previously and denied any recent trauma. His height was 177 cm, body weight was 59 kg, and his body mass index (BMI) was 18.8 kg/m². His blood pressure was 110/84 mmHg, heart rate was 70 beats/minute, respiratory rate 24/minute, body temperature 36.6°C, and right breath sounds were diminished on auscultation. Chest roentgenogram showed right pneumothorax without mediastinal shift (Figure 1). Chest CT showed subpleural bulla over left apical segments were found (Figure 2).



Figure 1. Chest roentgenogram showed right pneumothorax without mediastinal shift.

* E-mail: hirosato@md.tsukuba.ac.jp




Figure 2. Chest CT showed subpleural bulla over left apical segments were found.

The patient had undergone VATS for right pneumothorax. Subpleural bullae over the apical segment were excised with endoscopic linear staples. The patient was discharged on the sixth postoperative day. Five months after the surgery, the patient was seen at our hospital for the sudden onset of left chest pain. Upon examination, he was neither cyanotic nor tachypneic, but left hemithoraces was resonant to percussion. Chest roentgenogram showed left pneumothorax (Figure 3). VATS bullae resection with pleural abrasion was performed on the left hemothorax. He was discharged 4 days later and was followed up for 1 month at our outpatient clinic without recurrence.



Figure 3. Chest roentgenogram showed left pneumothorax.

3. Discussion

In urban areas, the emergency service must always be Spontaneous pneumothorax is most common in adolescents and young adults. Due to its tendency to recur, spontaneous pneumothorax is still troublesome for patients. Bullous lesions in the apex are evaluated as risk factors [7]. According to previous studies, the recurrence rate would be 16% to 50% if there was no surgical intervention [1,8-12]. With surgical treatment, its recurrence rate becomes less than 5% [13]. However, contralateral recurrence may also take place [2-6]. Sihoe et al. reported that 53% of patients with spontaneous pneumothorax were found to have contralateral bullous lesions, and it has been identified as one of the most important risk factors in contralateral pneumothorax [5]. Mitlehner et al. reviewed the value of CT scan in detecting bullae and bleb formation of the lung in 35 patients with spontaneous pneumothorax. CT scans showed pathological lung changes in 31 of 35 patients. Contralateral bullae and blebs were found in 23 of them. However, no correlation between recurrences and anatomical status such as number, size and distribution of blebs/bullae was found [14]. In addition, some researchers indicated that lower body mass index (BMI) was also a risk factor for bilateral and contralateral pneumothorax [15-17]. Sadikot et al. reported that recurrence was more common in taller men and in women [18]. Huang et al. recently reported the spontaneous bilateral and contralateral pneumothorax was significantly more frequent in patients with a lower BMI [16]. They also reported that BMI lower than BMI < 18.5 kg/m² was indicators for patients with spontaneous pneumothorax to receive single-stage bilateral surgery [16]. In our patient, BMI was over this level of < 18.5 kg/m², but we must not underestimate the fact that his BMI was low nonetheless. For spontaneous pneumothorax that could not be treated by less invasive procedures such as observation, manual aspiration, drainage or thoracoscopy, surgical treatment has been the last therapeutic resort. For recurrent ipsilateral or bilateral pneumothorax, or pneumothorax with persistent air leak, bilateral synchronous pneumothorax, it has been widely accepted that surgery is the treatment of choice. In a review of the management of spontaneous pneumothorax Tschopp et al. reported that simultaneous bilateral pneumothorax and contralateral recurrence are quite often indications for surgical treatment of both sides [19]. According to the guideline for the management of spontaneous pneumothorax by British Thoracic Society, contralateral pneumothorax is also one of the accepted indications for operative in-

tervention [7]. Its recommendation issues from both the statistical findings and perceived risk of recurrence [7]. Recently Chou *et al.* showed that the preemptive VATS for the contralateral blebs/bullae effectively prevented the contralateral occurrence [20]. In our patient, single-stage surgery was not performed at the first presentation, as the patient did not want to receive VATS in both sides at that time. We underestimated the risk of contralateral pneumothorax because of the size and shape of the bullae that were usually observed. There has been no reports that evaluate what kinds of bullae will develop in pneumothorax. It is expected such studies will be carried out in future. In this case, we referred the patient to the

thoracic surgeons soon after the diagnosis of contralateral pneumothorax. This single-stage surgery will be one of choices of the common therapies, but it has not yet been evaluated as the standard one.

It is important to evaluate the existence of bullous lesion in both sides of the lungs at the time of the first presentation of pneumothorax, as a bullous lesion on the contralateral side on the chest CT scan is an independent risk factor for contralateral recurrence. Either chest physicians or thoracic surgeons must follow up with recognition of increased risk of contralateral pneumothorax onset in patients with bullous lesions on the contralateral side as well as low BMI.

References

- [1] Schramel FM, Postmus PE, Vanderschueren RG. Current aspects of spontaneous pneumothorax. *Eur Respir J* 1997; 10: 1372-1379
- [2] Ikeda M, Uno A, Yamane Y, *et al.* Median sternotomy with bilateral bullous resection for unilateral spontaneous pneumothorax, with special reference to operative indications. *J Thorac Cardiovasc Surg* 1988; 96: 615-620
- [3] Gamondes JP, Wiesendanger T, Bouvier H, *et al.* Recurrent spontaneous pneumothorax in young subjects: treatment by one-stage bilateral apical pleurectomy through the axillary route. *Presse Med* 1987; 16: 423-426
- [4] Driscoll PJ, Aronstam EM. Experiences in the management of recurrent spontaneous pneumothorax. *J Thorac Cardiovasc Surg* 1961;42:174-178
- [5] Sihoe AD, Yim AP, Lee TW, Wan S, Yuen EH, Wan IY, *et al.* Can CT scanning be used to select patients with unilateral primary spontaneous pneumothorax for bilateral surgery? *Chest* 2000; 118: 380-383
- [6] Ouanes-Besbes L, Golli M, Knani J, Dachraoui F, Nciri N, El Atrous S, *et al.* Prediction of recurrent spontaneous pneumothorax: CT scan findings versus management features. *Respir Med* 2007;101:230-236
- [7] MacDuff A, Arnold A, Harvey J; BTS Pleural Disease Guideline Group. Management of spontaneous pneumothorax: British Thoracic Society Pleural Disease Guideline 2010. *Thorax* 2010;65 Suppl 2:ii18-31
- [8] Guo Y, Xie C, Rodriguez RM, Light RW. Factors related to recurrence of spontaneous pneumothorax. *Respirology* 2005; 10: 378-384
- [9] Baumann MH, Strange C. Treatment of spontaneous pneumothorax: a more aggressive approach? *Chest* 1997; 112: 789-804
- [10] Baumann MH, Noppen M. Pneumothorax. *Respirology* 2004; 9: 157-164
- [11] Chan SS, Lam PK. Simple aspiratory as initial treatment for primary spontaneous pneumothorax: results of 91 consecutive cases. *J Emerg Med* 2005; 28: 133-138
- [12] Ng CS, Lee TW, Wan S, Yim AP. Video-assisted thoracic surgery in the management of spontaneous pneumothorax: the current status. *Postgrad Med J* 2006; 82: 179-185
- [13] Weissberg D, Refaely Y. Pneumothorax: experience with 1,199 patients. *Chest* 2000; 117: 1279-1285
- [14] Mittlehner W, Friedrich M, Dissmann W. Value of computer tomography in the detection of bullae and blebs in patients with primary spontaneous pneumothorax. *Respiration* 1992; 59: 221-227
- [15] Huang TW, Lee SC, Cheng YL, Tzao C, Hsu HH, Chang H, Chen JC. Contralateral recurrence of primary spontaneous pneumothorax. *Chest* 2007; 132: 1146-1150
- [16] Huang TW, Cheng YL, Tzao C, Hung C, Hsu HH, Chen JC, Lee SC. Factors related to primary bilateral spontaneous pneumothorax. *Thorac Cardiovasc Surg* 2007; 55: 310-312
- [17] Ayed AK, Bazerbashi S, Ben-Nakhi M, *et al.* Risk factors of spontaneous pneumothorax in Kuwait. *Med Princ Pract* 2006; 15: 338-342
- [18] Sadikot RT, Greene T, Meadows K, Arnold AG. Recurrence of primary spontaneous pneumothorax. *Thorax* 1997; 52: 805-809
- [19] Tschopp JM, Rami-Porta R, Noppen M, Astoul P. Management of spontaneous pneumothorax: state of the art. *Eur Respir J* 2006; 28: 637-650
- [20] Chou SH, Li HP, Lee JY, Chang SJ, Lee YL, Chang YT, Kao EL, Dai ZK, Huang MF. Is prophylactic treatment of contralateral blebs in patients with primary spontaneous pneumothorax indicated? *J Thorac Cardiovasc Surg* 2010; 139: 1241-1245