

## RESEARCH PAPER

### SPECTRAL APPROXIMATIONS TO THE FRACTIONAL INTEGRAL AND DERIVATIVE

Changpin Li <sup>1</sup>, Fanhai Zeng <sup>2</sup>, Fawang Liu <sup>3</sup>

#### Abstract

In this paper, the spectral approximations are used to compute the fractional integral and the Caputo derivative. The effective recursive formulae based on the Legendre, Chebyshev and Jacobi polynomials are developed to approximate the fractional integral. And the succinct scheme for approximating the Caputo derivative is also derived. The collocation method is proposed to solve the fractional initial value problems and boundary value problems. Numerical examples are also provided to illustrate the effectiveness of the derived methods.

*Editorial Note:* The first and third authors of this paper have recently received prestigious awards at the 5th International Symposium “Fractional Differentiation and Applications’ 2012” in China, May 14-17, 2012. Professor Changpin Li, together with his PhD student Yutian Ma, received the “Riemann-Liouville Award” for Best FDA Paper (theory), FDA #077: Fractional Dynamical System and Its Linearization Theorem. Professor Fawang Liu got the “Mittag-Leffler Award” which is given once per two years for a cumulative contribution to the field of Fractional Derivative and Applications, see details at <http://em.hhu.edu.cn/fda12/Awards.html>.

*MSC 2010:* 26A33 (main), 65L12, 15A99, 39A70

*Key Words and Phrases:* fractional integral, Caputo derivative, spectral approximation, Jacobi polynomials

## 1. Introduction

Fractional calculus (including the operators of fractional order integration and differentiation) is as old as its more familiar integer-order counterpart, the classical Calculus, and has a long history, see e.g. [15, 22, 27], etc. Since then, fractional calculus has undergone a rapid development in theoretical aspects and recently has been found wide applications in many areas of science and engineering, for example [3, 9, 11, 12, 16, 17, 20, 22, 28, 30, 32].

The operator of fractional derivative is more complicated than the classical one, and its calculation is also more difficult than in the integer order case. In [4], numerical algorithms for establishing the fractional integral and the Caputo derivative were considered. Li et al. [10] also developed numerical algorithms based on the piecewise polynomial interpolation to approximate the fractional integral and the Caputo derivative, and to solve fractional differential equations. An automatic quadrature method based on the Chebyshev polynomials was presented for approximating the Caputo derivative in [26]. Some other computational schemes, such as the L1, L2 and L2C schemes, etc., are also introduced [8, 11, 13, 14, 16, 18, 19, 21, 23, 25, 28, 29, 31].

To increase the calculation accuracy, spectral approximations are often chosen. In this paper, we derive effective algorithms to approximate the fractional integral by using the Legendre, Chebyshev and Jacobi polynomials. A numerical algorithm to calculate the Caputo derivative is also derived based on the above computational schemes. Besides, we propose a kind of fractional operational matrix, which can be seen as a generalization of the classical derivative. When the fractional order of the Caputo derivative reduces to an integer, the derived fractional operational matrix reduces to the classical differential matrix. Applications of the constructed algorithms are illustrated to compute the fractional integral, Caputo derivative and to solve fractional ordinary differential equations. Numerical experiments are described to verify the proposed numerical algorithms.

The remainder of this paper is organized as follows. In Section 2, we introduce several definitions of fractional calculus' operators and the Legendre, Chebyshev and Jacobi polynomials. Numerical algorithms for calculating the fractional integral and the Caputo derivative are presented in Sections 3 and 4, respectively. The applications of the methods are illustrated in Section 5. Numerical examples are presented in Section 6, and the conclusion is included in the last section.

## 2. Definitions and notations

In this section we introduce the definitions of the operators of fractional calculus and of the Legendre, Chebyshev and Jacobi polynomials, used later on.

**DEFINITION 2.1.** The fractional integral (or the Riemann–Liouville integral) of order  $\alpha > 0$  of a given function  $f(t)$  is defined as

$$D_{a,t}^{-\alpha} f(t) = \frac{1}{\Gamma(\alpha)} \int_a^t (t-s)^{\alpha-1} f(s) \, ds, \quad (2.1)$$

where  $\Gamma(\cdot)$  is the Euler's gamma function.

There exist several kinds of definitions of fractional derivatives. However, in practical applications as engineering, physics etc., the Caputo derivative is mostly used, as introduced below.

**DEFINITION 2.2.** The Caputo derivative of order  $\alpha > 0$  of a given function  $f(t)$  is defined as

$$\begin{aligned} {}_C D_{a,t}^{\alpha} f(t) &= D_{a,t}^{-(n-\alpha)} \left[ f^{(n)}(t) \right] \\ &= \frac{1}{\Gamma(n-\alpha)} \int_a^t (t-s)^{n-\alpha-1} f^{(n)}(s) \, ds, \end{aligned} \quad (2.2)$$

where  $n$  is a positive integer and  $n-1 < \alpha \leq n$ .

Next, we introduce the Legendre, Chebyshev and Jacobi polynomials.

The Legendre polynomials  $\{L_j(x)\}$ ,  $x \in [-1, 1]$  satisfy the three-term recurrence relation

$$\begin{aligned} L_0(x) &= 1, \quad L_1(x) = x, \\ L_{j+1} &= \frac{2j+1}{j+1} x L_j(x) - \frac{j}{j+1} L_{j-1}(x), \quad j \geq 1. \end{aligned} \quad (2.3)$$

Their properties which will be used later on are listed as follows:

$$(2j+1)L_j(x) = L'_{j+1}(x) - L'_{j-1}(x), \quad j \geq 1, \quad (2.4)$$

$$L_j(\pm 1) = (\pm 1)^j, \quad L'_j(\pm 1) = \frac{1}{2}(\pm 1)^{j-1} j(j+1). \quad (2.5)$$

The Chebyshev polynomials  $\{T_j(x)\}, x \in [-1, 1]$  satisfy the three-term recurrence relation

$$\begin{aligned} T_0(x) &= 1, \quad T_1(x) = x, \\ T_{j+1}(x) &= 2xT_j(x) - T_{j-1}(x), \quad j \geq 1. \end{aligned} \quad (2.6)$$

Their fundamental properties are presented below:

$$2T_j(x) = \frac{1}{j+1}T'_{j+1}(x) - \frac{1}{j-1}T'_{j-1}(x), \quad j \geq 2, \quad (2.7)$$

$$T_j(\pm 1) = (\pm 1)^j, \quad T'_j(\pm 1) = \frac{1}{2}(\pm 1)^{j-1}j^2. \quad (2.8)$$

The Jacobi polynomials  $\{P_j^{a,b}(x)\}, a, b > -1, x \in [-1, 1]$  are given by the following three-term recurrence relation ([24]):

$$\begin{aligned} P_0^{a,b}(x) &= 1, \quad P_1^{a,b}(x) = \frac{1}{2}(a+b+2)x + \frac{1}{2}(a-b), \\ P_{j+1}^{a,b}(x) &= (A_j^{a,b}x - B_j^{a,b})P_j^{a,b}(x) - C_j^{a,b}P_{j-1}^{a,b}(x), \quad n \geq 1, \end{aligned} \quad (2.9)$$

where

$$\begin{aligned} A_j^{a,b} &= \frac{(2j+a+b+1)(2j+a+b+2)}{2(j+1)(j+a+b+1)}, \\ B_j^{a,b} &= \frac{(b^2-a^2)(2j+a+b+1)}{2(j+1)(j+a+b+1)(2j+a+b)}, \\ C_j^{a,b} &= \frac{(j+a)(j+b)(2j+a+b+2)}{(j+1)(j+a+b+1)(2j+a+b)}. \end{aligned} \quad (2.10)$$

Here, we list some useful properties of the Jacobi polynomials that will be used in the present paper [24]:

$$P_j^{a,b}(1) = \binom{j+a}{j} = \frac{\Gamma(j+a+1)}{j!\Gamma(a+1)}, \quad P_j^{a,b}(-1) = (-1)^j \frac{\Gamma(j+b+1)}{j!\Gamma(b+1)}. \quad (2.11)$$

$$\frac{d^m}{dx^m} P_j^{a,b}(x) = d_{j,m}^{a,b} P_{j-m}^{a+m,b+m}(x), \quad j \geq m, m \in \mathbb{N}, \quad (2.12)$$

where

$$d_{j,m}^{a,b} = \frac{\Gamma(j+m+a+b+1)}{2^m \Gamma(j+a+b+2)}. \quad (2.13)$$

$$P_j^{a,b}(x) = \hat{A}_j^{a,b} \frac{d}{dx} P_{j-1}^{a,b}(x) + \hat{B}_j^{a,b} \frac{d}{dx} P_j^{a,b}(x) + \hat{C}_j^{a,b} \frac{d}{dx} P_{j+1}^{a,b}(x), \quad j \geq 1, \quad (2.14)$$

in which

$$\begin{aligned}
\hat{A}_j^{a,b} &= \frac{-2(j+a)(j+b)}{(j+a+b)(2j+a+b)(2j+a+b+1)}, \\
\hat{B}_j^{a,b} &= \frac{2(a-b)}{(2j+a+b)(2j+a+b+2)}, \\
\hat{C}_j^{a,b} &= \frac{2(j+a+b+1)}{(2j+a+b+1)(2j+a+b+2)}.
\end{aligned} \tag{2.15}$$

If  $j = 1$ ,  $\hat{A}_1^{a,b}$  in (2.15) is set to be zero.

**REMARK 2.1.** The case  $a = b = 0$  in (2.9) yields the Legendre polynomials ( $P_j^{0,0}(x) = L_j(x)$ ). If  $a = b = -\frac{1}{2}$ , then  $P_j^{-\frac{1}{2}, -\frac{1}{2}}(x) = \frac{\Gamma(j+1/2)}{j! \sqrt{\pi}} T_j(x)$ ,  $j \geq 0$ .

### 3. Approximation to the fractional integral

In this section we develop algorithms to approximate the fractional integral of a given function. Among the three mentioned kinds of polynomials, the Legendre polynomials are commonly used. In this section we mainly study in details the Legendre approximation to the fractional integral. The Chebyshev and Jacobi approximations are almost the same as the Legendre approximation.

Let  $u(x)$  be a function defined on the interval  $[-1, 1]$ , and  $N$  be a positive integer. Denote by

$$p_N(x) = \sum_{j=0}^N \tilde{l}_j L_j(x), \tag{3.1}$$

where  $p_N(x)$  is an approximation of  $u(x)$ , and  $\tilde{l}_j$  are the coefficients determined by  $u(x)$ . If  $p_N(x)$  is an orthogonal projection of  $u(x)$ , then  $\tilde{l}_j$  can be determined by the orthogonality of  $\{L_j(x)\}$ . In this paper, we assume that  $p_N(x)$  is the interpolation of  $u(x)$ . If  $p_N(x)$  is the interpolation of  $u(x)$  on the Legendre–Gauss–Lobatto points  $\{x_k\}_{k=0}^N$ , then  $\tilde{l}_j$  can be determined by

$$\tilde{l}_j = \frac{1}{\gamma_j} \sum_{k=0}^N u(x_k) L_j(x_k) \omega_k, \tag{3.2}$$

where  $\gamma_j = \frac{2}{2j+1}$  for  $0 \leq j \leq N-1$ ,  $\gamma_N = \frac{2}{N}$ , and  $\{\omega_k\}_{k=0}^N$  are the corresponding quadrature weights [24].

Therefore, for any  $\alpha > 0$ , the fractional integral  $D_{-1,x}^{-\alpha} u(x)$  can be approximated by

$$\begin{aligned}
D_{-1,x}^{-\alpha} u(x) &\approx D_{-1,x}^{-\alpha} p_N(x) = \frac{1}{\Gamma(\alpha)} \int_{-1}^x (x-s)^{\alpha-1} p_N(s) ds \\
&= \frac{1}{\Gamma(\alpha)} \sum_{j=0}^N \tilde{l}_j \int_{-1}^x (x-s)^{\alpha-1} L_j(s) ds = \sum_{j=0}^N \tilde{l}_j \hat{L}_j^{\alpha}(x),
\end{aligned} \tag{3.3}$$

where  $\hat{L}_j^{\alpha}(x) = \frac{1}{\Gamma(\alpha)} \int_{-1}^x (x-s)^{\alpha-1} L_j(s) ds$ . Next, we give an effective recurrence formula to calculate  $\hat{L}_j^{\alpha}(x)$ .

We can easily get

$$\hat{L}_0^{\alpha}(x) = \frac{(x+1)^{\alpha}}{\Gamma(\alpha+1)} \quad \text{and} \quad \hat{L}_1^{\alpha}(x) = \frac{x(x+1)^{\alpha}}{\Gamma(\alpha+1)} - \frac{\alpha(x+1)^{\alpha+1}}{\Gamma(\alpha+2)}$$

from (2.3). For  $j \geq 1$ , by using (2.3), we have

$$\begin{aligned}
\hat{L}_{j+1}^{\alpha}(x) &= \frac{1}{\Gamma(\alpha)} \int_{-1}^x (x-s)^{\alpha-1} L_{j+1}(s) ds \\
&= \frac{1}{\Gamma(\alpha)} \frac{1}{j+1} \int_{-1}^x (x-s)^{\alpha-1} \left[ (2j+1)sL_j(s) - jL_{j-1}(s) \right] ds \\
&= \frac{1}{j+1} \left\{ (2j+1)x\hat{L}_j^{\alpha}(x) - j\hat{L}_{j-1}^{\alpha}(x) - \frac{2j+1}{\Gamma(\alpha)} \int_{-1}^x (x-s)^{\alpha} L_j(s) ds \right\}.
\end{aligned} \tag{3.4}$$

Noticing that  $(2j+1)L_j(x) = L'_{j+1}(x) - L'_{j-1}(x)$  for  $j \geq 1$  (see Eq. (2.4)), we have

$$\begin{aligned}
\hat{L}_{j+1}^{\alpha}(x) &= \frac{1}{j+1} \left\{ (2j+1)x\hat{L}_j^{\alpha}(x) - j\hat{L}_{j-1}^{\alpha}(x) \right. \\
&\quad \left. - \frac{1}{\Gamma(\alpha)} \int_{-1}^x (x-s)^{\alpha} (L'_{j+1}(s) - L'_{j-1}(s)) ds \right\} \\
&= \frac{1}{j+1} \left\{ (2j+1)x\hat{L}_j^{\alpha}(x) - j\hat{L}_{j-1}^{\alpha}(x) \right. \\
&\quad \left. - \frac{1}{\Gamma(\alpha)} \left[ (x-s)^{\alpha} (L_{j+1}(s) - L_{j-1}(s)) \right]_{-1}^x - \alpha(\hat{L}_{j+1}^{\alpha}(x) - \hat{L}_{j-1}^{\alpha}(x)) \right\} \\
&= \frac{1}{j+1} \left\{ (2j+1)x\hat{L}_j^{\alpha}(x) - j\hat{L}_{j-1}^{\alpha}(x) - \alpha(\hat{L}_{j+1}^{\alpha}(x) - \hat{L}_{j-1}^{\alpha}(x)) \right\}.
\end{aligned} \tag{3.5}$$

Hence, for  $j \geq 1$ , we get the following recurrence relation

$$\hat{L}_{j+1}^{\alpha}(x) = \frac{1}{j+1+\alpha} \left\{ (2j+1)x\hat{L}_j^{\alpha}(x) - (j-\alpha)\hat{L}_{j-1}^{\alpha}(x) \right\}. \tag{3.6}$$

So,  $\hat{L}_j^{\alpha}(x)$  can be calculated by the following formula

$$\left\{ \begin{array}{l} \widehat{L}_0^\alpha(x) = \frac{(x+1)^\alpha}{\Gamma(\alpha+1)}, \\ \widehat{L}_1^\alpha(x) = \frac{x(x+1)^\alpha}{\Gamma(\alpha+1)} - \frac{\alpha(x+1)^{\alpha+1}}{\Gamma(\alpha+2)}, \\ \widehat{L}_{j+1}^\alpha(x) = \frac{1}{j+1+\alpha} \left\{ (2j+1)x\widehat{L}_j^\alpha(x) - (j-\alpha)\widehat{L}_{j-1}^\alpha(x) \right\}, \quad j \geq 1. \end{array} \right. \quad (3.7)$$

Therefore,  $D_{-1,x}^{-\alpha}u(x)$  is approximated by

$$D_{-1,x}^{-\alpha}u(x) \approx D_{-1,x}^{-\alpha}p_N(x) = \sum_{j=0}^N \tilde{l}_j \widehat{L}_j^\alpha(x),$$

where  $\widehat{L}_j^\alpha(x)$  is defined by (3.7), and  $\tilde{l}_j$  is defined by (3.2).

Similarly, we can get the similar results for the Chebyshev polynomials and the Jacobi polynomials.

For the Chebyshev polynomials, let  $\widehat{T}_j^\alpha(x) = \frac{1}{\Gamma(\alpha)} \int_{-1}^x (x-s)^{\alpha-1} T_j(s) ds$ , by many calculations, one can get

$$\left\{ \begin{array}{l} \widehat{T}_0^\alpha(x) = \frac{(x+1)^\alpha}{\Gamma(\alpha+1)}, \\ \widehat{T}_1^\alpha(x) = \frac{x(x+1)^\alpha}{\Gamma(\alpha+1)} - \frac{\alpha(x+1)^{\alpha+1}}{\Gamma(\alpha+2)}, \\ \widehat{T}_2^\alpha(x) = \frac{4x}{2+\alpha} \widehat{T}_1^\alpha(x) - \frac{2}{2+\alpha} \widehat{T}_0^\alpha(x) + \frac{\alpha(x+1)^\alpha}{(2+\alpha)\Gamma(\alpha+1)}, \\ \widehat{T}_{j+1}^\alpha(x) = \frac{2(j+1)x}{j+1+\alpha} \widehat{T}_j^\alpha(x) - \frac{(j+1)(j-1-\alpha)}{(j+1+\alpha)(j-1)} \widehat{T}_{j-1}^\alpha(x) \\ \quad + \frac{2(-1)^j \alpha (x+1)^\alpha}{\Gamma(\alpha+1)(j+1+\alpha)(j-1)}, \quad j \geq 2. \end{array} \right. \quad (3.8)$$

If  $u(x) \approx p_N(x) = \sum_{j=0}^N \tilde{t}_j T_j(x)$ , then  ${}_CD_{-1,x}^{-\alpha}u(x)$  can be approximated by

$$D_{-1,x}^{-\alpha}u(x) \approx D_{-1,x}^{-\alpha}p_N(x) = \sum_{j=0}^N \tilde{t}_j \widehat{T}_j^\alpha(x),$$

where  $\tilde{t}_j$  can be similarly determined, see [24] for details.

For the Jacobi polynomials, we denote by

$$\widehat{P}_j^{a,b,\alpha}(x) = \frac{1}{\Gamma(\alpha)} \int_{-1}^x (x-s)^{\alpha-1} P_j^{a,b}(s) ds.$$

Using the recurrence formulae (2.9)–(2.10), the properties (2.11) and (2.14), and tedious calculations, we have

$$\left\{ \begin{array}{l} \hat{P}_0^{a,b,\alpha}(x) = \frac{(x+1)^\alpha}{\Gamma(\alpha+1)}, \\ \hat{P}_1^{a,b,\alpha}(x) = \frac{a+b+2}{2} \left( \frac{x(x+1)^\alpha}{\Gamma(\alpha+1)} - \frac{\alpha(x+1)^{\alpha+1}}{\Gamma(\alpha+2)} \right) + \frac{a-b}{2} \hat{P}_0^{a,b,\alpha}(x), \\ \hat{P}_{j+1}^{a,b,\alpha}(x) = \frac{A_j^{a,b}x - B_j^{a,b} - \alpha A_j^{a,b} \hat{B}_j^{a,b}}{1 + \alpha A_j^{a,b} \hat{C}_j^{a,b}} \hat{P}_j^{a,b,\alpha}(x) \\ \quad - \frac{C_j^{a,b} + \alpha A_j^{a,b} \hat{A}_j^{a,b}}{1 + \alpha A_j^{a,b} \hat{C}_j^{a,b}} \hat{P}_{j-1}^{a,b,\alpha}(x) \\ \quad + \frac{\alpha A_j^{a,b} \left( \hat{A}_j^{a,b} P_{j-1}^{a,b}(-1) + \hat{B}_j^{a,b} P_j^{a,b}(-1) + \hat{C}_j^{a,b} P_{j+1}^{a,b}(-1) \right)}{\Gamma(\alpha+1) \left( 1 + \alpha A_j^{a,b} \hat{C}_j^{a,b} \right)} (x+1)^\alpha, \\ j \geq 1. \end{array} \right. \quad (3.9)$$

If  $u(x) \approx p_N(x) = \sum_{j=0}^N \tilde{p}_j^{a,b} P_j^{a,b}(x)$ , then  $D_{-1,x}^{-\alpha} u(x)$  can be approximated by

$$D_{-1,x}^{-\alpha} u(x) \approx D_{-1,x}^{-\alpha} p_N(x) = \sum_{j=0}^N \tilde{p}_j^{a,b} \hat{P}_j^{a,b,\alpha}(x),$$

where  $\tilde{p}_j^{a,b}$  can be determined almost similarly as  $\tilde{l}_j$  in (3.2).

**REMARK 3.1.** If  $a = b = 0$ , then  $\hat{P}_j^{0,0,\alpha}(x) = \hat{L}_j^\alpha(x)$ . If  $a = b = -\frac{1}{2}$ , then  $\hat{P}_j^{-\frac{1}{2},-\frac{1}{2},\alpha}(x) = \frac{\Gamma(j+1/2)}{j!\sqrt{\pi}} \hat{T}_j^\alpha(x)$ .

#### 4. Approximation to the Caputo derivative

This section deals with the numerical approximation of the Caputo derivative of a given function  $u(x)$ ,  $x \in [-1, 1]$ . The algorithm is based on the numerical approximation of the fractional integral derived in the previous section. Among the Legendre, Chebyshev and Jacobi polynomials, the Jacobi polynomials are the most general. We study the Jacobi approximation to the Caputo derivative.

Suppose that  $p_N(x)$  is the approximate polynomial of  $u(x)$ , which can be expressed as

$$p_N(x) = \sum_{j=0}^N \tilde{p}_j^{a,b} P_j^{a,b}(x), \quad x \in [-1, 1]. \quad (4.1)$$



Let  $n - 1 < \alpha < n$ ,  $n$  is a positive integer, and the  $\alpha$ th Caputo derivative of  $p_N(x)$  reads as

$$\begin{aligned} {}_C D_{-1,x}^\alpha p_N(x) &= \frac{1}{\Gamma(n-\alpha)} \int_{-1}^x (x-s)^{n-\alpha-1} \frac{d^n}{ds^n} p_N(s) ds \\ &= \frac{1}{\Gamma(n-\alpha)} \sum_{j=0}^N \tilde{p}_j^{a,b} \int_{-1}^x (x-s)^{n-\alpha-1} \frac{d^n}{ds^n} P_j^{a,b}(s) ds \\ &= \sum_{j=n}^N \tilde{p}_j^{a,b} d_{j,n}^{a,b} \left( \frac{1}{\Gamma(n-\alpha)} \int_{-1}^x (x-s)^{n-\alpha-1} P_j^{a+n,b+n}(s) ds \right) \\ &= \sum_{j=n}^N \tilde{p}_j^{a,b} d_{j,n}^{a,b} \hat{P}_j^{a+n,b+n,n-\alpha}(x), \end{aligned}$$

where Eq. (2.12) is used.  $d_{j,n}^{a,b}$  is defined by (2.13), and  $\hat{P}_j^{a+n,b+n,n-\alpha}(x)$  is defined by (3.9).

Denote by

$$D_j^{a,b,\alpha,m} = \frac{1}{\Gamma(n-\alpha)} \int_{-1}^x (x-s)^{n-\alpha-1} \frac{d^m}{ds^m} P_j^{a,b}(s) ds, \quad 0 \leq m \leq n,$$

one has

$$D_j^{a,b,\alpha,n}(x) = d_{j,n}^{a,b} \hat{P}_j^{a+n,b+n,n-\alpha}(x), \quad (4.2)$$

where  $D_j^{a,b,\alpha,n}(x) = 0$  for  $0 \leq j \leq n-1$ ,  $d_{j,n}^{a,b}$  is given by (2.13), and  $\hat{P}_j^{a+n,b+n,n-\alpha}$  is defined by (3.9).

Therefore, the  $\alpha$ th Caputo derivative  ${}_C D_{-1,x}^\alpha u(x)$  can be approximated by

$${}_C D_{-1,x}^\alpha u(x) \approx {}_C D_{-1,x}^\alpha p_N(x) = \sum_{j=n}^N \tilde{p}_j^{a,b} D_j^{a,b,\alpha,n}(x),$$

where  $D_j^{a,b,\alpha,n}(x)$  is given by (4.2) and  $\tilde{p}_j^{a,b}$  can be similarly obtained as  $\tilde{l}_j$  in (3.2).

## 5. Applications of the algorithms

Let  $u(x)$  be a real-valued function defined on the interval  $[x_a, x_b]$ , and  $\hat{x}_i$  ( $i = 0, 1, \dots, N$ ) be the collocation points on the reference interval  $[-1, 1]$ , then  $x_i = \frac{1}{2}[(x_b - x_a)\hat{x}_i + x_a + x_b]$  are the corresponding collocation points on  $[x_a, x_b]$ . Similarly to the previous section, we use the more general Jacobi polynomials, to illustrate our numerical methods. For a positive integer  $N$ ,

suppose that  $u(x)$  can be approximated by the following polynomial

$$u(x) \approx p_N(x) = \sum_{j=0}^N \tilde{p}_j^{a,b} P_j^{a,b}(\hat{x}), \quad \hat{x} = \frac{2x - x_a - x_b}{x_b - x_a} \in [-1, 1], \quad (5.1)$$

Therefore, for any  $\alpha > 0$ , we have

$$\begin{aligned} D_{x_a, x}^{-\alpha} p_N(x) &= \frac{1}{\Gamma(\alpha)} \sum_{j=0}^N \tilde{p}_j^{a,b} D_{x_a, x}^{-\alpha} P_j^{a,b}(\hat{x}) \\ &= \left( \frac{x_b - x_a}{2} \right)^\alpha \sum_{j=0}^N \tilde{p}_j^{a,b} \hat{P}_j^{a,b, \alpha}(\hat{x}), \end{aligned} \quad (5.2)$$

in which

$$\begin{aligned} \frac{1}{\Gamma(\alpha)} D_{x_a, x}^{-\alpha} P_j^{a,b}(\hat{x}) &= \frac{1}{\Gamma(\alpha)} \int_{x_a}^x (x-s)^{\alpha-1} P_j^{a,b}(\hat{s}) ds \\ &= \left( \frac{x_b - x_a}{2} \right)^\alpha \frac{1}{\Gamma(\alpha)} \int_{-1}^{\hat{x}} (\hat{x} - \hat{s})^{\alpha-1} P_j^{a,b}(\hat{s}) d\hat{s} \\ &= \left( \frac{x_b - x_a}{2} \right)^\alpha \hat{P}_j^{a,b, \alpha}(\hat{x}). \end{aligned} \quad (5.3)$$

Here  $s = \frac{1}{2}[(x_b - x_a)\hat{s} + x_a + x_b]$  and  $x = \frac{1}{2}[(x_b - x_a)\hat{x} + x_a + x_b]$ ,  $\hat{s}, \hat{x} \in [-1, 1]$  are used.  $\hat{P}_j^{a,b, \alpha}(\hat{x})$  is defined by (3.9).

So, the fractional integral of  $u(x)$  at any point  $x \in [x_a, x_b]$  can be approximated by (5.2). If we choose  $x = x_i$  ( $i = 0, 1, \dots, N$ ), then we have

$$\begin{pmatrix} D_{x_a, x_0}^{-\alpha} u(x_0) \\ D_{x_a, x_1}^{-\alpha} u(x_1) \\ \vdots \\ D_{x_a, x_N}^{-\alpha} u(x_N) \end{pmatrix} \approx \begin{pmatrix} D_{x_a, x_0}^{-\alpha} p_N(x_0) \\ D_{x_a, x_1}^{-\alpha} p_N(x_1) \\ \vdots \\ D_{x_a, x_N}^{-\alpha} p_N(x_N) \end{pmatrix} = {}_C D_{x_a, x_b}^{(a,b, -\alpha)} \tilde{\mathbf{p}}^{a,b}, \quad (5.4)$$

where  $\tilde{\mathbf{p}}^{a,b} = (\tilde{p}_0^{a,b}, \tilde{p}_1^{a,b}, \dots, \tilde{p}_N^{a,b})^T$ , and

$$\left[ D_{x_a, x_b}^{(a,b, -\alpha)} \right]_{i,j} = \left( \frac{x_b - x_a}{2} \right)^\alpha \hat{P}_j^{a,b, \alpha}(\hat{x}_i), \quad i, j = 0, \dots, N.$$

Similarly, for  $n-1 < \alpha < n \in \mathbb{N}$ , we can also get the following formula for the Caputo derivative

$$\begin{pmatrix} D_{x_a, x_0}^{\alpha} u(x_0) \\ D_{x_a, x_1}^{\alpha} u(x_1) \\ \vdots \\ D_{x_a, x_N}^{\alpha} u(x_N) \end{pmatrix} \approx \begin{pmatrix} D_{x_a, x_0}^{\alpha} p_N(x_0) \\ D_{x_a, x_1}^{\alpha} p_N(x_1) \\ \vdots \\ D_{x_a, x_N}^{\alpha} p_N(x_N) \end{pmatrix} = D_{x_a, x_b}^{(a,b, \alpha)} \tilde{\mathbf{p}}^{a,b}, \quad (5.5)$$

where

$$\left[ {}_C D_{x_a, x_b}^{(a, b, \alpha)} \right]_{i, j} = \left( \frac{x_b - x_a}{2} \right)^{-\alpha} D_j^{a, b, \alpha, n}(\hat{x}_i), \quad i, j = 0, \dots, N, \quad (5.6)$$

and  $D_j^{a, b, \alpha, n}$  is defined by (4.2).

**REMARK 5.1.** For a given positive integer  $N$ , the matrix  ${}_C D_{x_a, x_b}^{(a, b, \alpha)}$  can be calculated effectively by  $O(N^2)$  arithmetic operations. The similar work can be found in [21], where the operational matrix  $\mathbf{D}^{(\alpha)}$  based on the explicit form of the Legendre polynomials was obtained, which takes  $O(N^3)$  arithmetic operations. The similar matrix  $\mathbf{D}^{(\alpha)}$  based on the Chebyshev polynomials can be found in [5, 6], where the computational complexity is  $O(N^3)$ .

**REMARK 5.2.** If  $p_N(x)$  (see (5.1)) is the Legendre-Gauss-Lobatto interpolation of  $u(x)$ ,  $u \in H^r(I)$ ,  $I = [x_a, x_b]$ , then

$$\|u - p_N\|_{L_\infty(I)} \leq C N^{3/4-r} \|u\|_{H^r(I)}, \quad r \in \mathbb{N},$$

where  $C$  is a positive constant (see [2]). Therefore, we can get the following the error bounds

$$\|D_{x_a, x}^{-\alpha}(u - p_N)\|_{L_\infty(I)} \leq C N^{3/4-r} \|u\|_{H^r(I)}, \quad \alpha > 0, r \geq 1,$$

and

$$\|{}_C D_{x_a, x}^\alpha(u - p_N)\|_{L_\infty(I)} \leq C N^{3/4+2n-r} \|u\|_{H^r(I)}, \quad n-1 < \alpha < n, n \in \mathbb{N}, r \geq 2n.$$

The above estimates will be verified by the numerical experiments in the following section.

Next we consider the eigenvalues of the Caputo fractional operator. Let  $x_a = -1, x_b = 1$ , and  $x_i (i = 0, 1, \dots, N)$  be the Legendre-Gauss-Lobatto points. We get the matrix  ${}_C D_{-1, 1}^{(0, 0, \alpha)} \in R^{(N+1) \times (N+1)}$  from Eq. (5.6).  ${}_C D_{-1, 1}^{(0, 0, \alpha)}$  is actually a kind of the operational differential matrix corresponding to the Caputo derivative. If  $\alpha = n$ , i.e., a positive integer, then  ${}_C D_{-1, 1}^{(0, 0, \alpha)}$  reduces to the classical differential matrix  $D^{(n)}$ .

In the following, we study numerically the spectral radius of the Caputo derivative operator. Consider the following model problem

$$\begin{cases} {}_C D_{-1, x}^\alpha u(x) = \lambda u(x), & x \in [-1, 1], \quad 0 < \alpha \leq 2, \\ u(-1) = u_{-1}, \quad u(1) = u_1, \end{cases} \quad (5.7)$$




FIGURE 1. The spectral radius associated with the Caputo fractional operator for  $0 < \alpha \leq 1$ .

where besides  $u(-1) = u_{-1}$ , the condition  $u(1) = u_1$  is necessary for  $1 < \alpha \leq 2$ .

For  $0 < \alpha \leq 1$ , let  $p_N(x)$  (see (5.1)) be the interpolation of  $u(x)$  on the Legendre–Gauss–Lobatto points  $\{x_i\}$ . Letting  $p_N(x)$  satisfy the equation (5.7) at  $x = x_i$  ( $i = 1, 2, \dots, N$ ), and combining the initial condition  $u(-1) = u_{-1}$ , we have

$$D\tilde{\mathbf{p}}^{a,b} = \lambda\tilde{\mathbf{p}}^{a,b}, \quad (5.8)$$

where  $[D]_{i,j} = [{}_CD_{-1,1}^{(0,0,\alpha)}]_{i,j}$  ( $i = 1, 2, \dots, N; j = 0, 1, \dots, N$ ), and  $[D]_{0,j} = (-1)^j$  ( $j = 0, 1, \dots, N$ ). It is easy to see that  $\lambda$  is just the eigenvalue of the matrix  $D$ . We know that, if  $\alpha = 1$ , the spectral radius  $\rho(D)$  of  $D$  satisfies the following relation

$$\rho(D) \leq C_0 N^2, \quad C_0 > 0.$$

For  $1 < \alpha \leq 2$ , we can also get almost the same relation as (5.8), except that the last row of  $D$  in (5.8) is replaced by  $[D]_{N,j} = 1$  ( $j = 0, 1, \dots, N$ ). We plot the spectral radius of  $D$  for different  $\alpha$  and  $N$  in Figures 1 and 2. From Figures 1 and 2, we can see that the spectral radius  $\rho(D)$  of  $D$  is bounded by

$$\rho(D) \leq C_0 N^{2\alpha}, \quad 0 < \alpha \leq 2, C_0 > 0.$$




FIGURE 2. The spectral radius associated with the Caputo fractional operator for  $1 < \alpha \leq 2$ .

Next, we simply illustrate how to use our method to solve the fractional differential equations. Consider the fractional equation in the following form

$$\begin{cases} A(x)u'(x) + B(x) {}_C D_{0,x}^\alpha u(x) + C(x)u(x) = f(x), & x \in (0, L], \\ u(0) = u_0, \end{cases} \quad (5.9)$$

where  $0 < \alpha < 1$ ,  $A(x)$ ,  $B(x)$  and  $C(x)$  are real-valued functions.

Suppose that  $p_N(x)$  (see Eq. (5.1) and set  $x_a = 0, x_b = L$ ) is the approximate solution of  $u(x)$ ,  $x_i$  ( $i = 0, 1, \dots, N$ ) are the collocation points on the interval  $[0, L]$  satisfying  $p_N(x_i) = u(x_i)$  ( $i = 0, 1, \dots, N$ ) and

$$A(x_i)p_N'(x_i) + B(x_i) {}_C D_{0,x_i}^\alpha p_N(x_i) + C(x_i)u(x_i) = f(x_i), \quad i = 1, 2, \dots, N.$$

Noting  $p_N(x_0) = u_0$ , we get the following algebraic equation

$$M\tilde{\mathbf{p}}^{a,b} = F, \quad (5.10)$$

where  $F = (u_0, f(x_1), f(x_2), \dots, f(x_N))^T$ , and

$$\begin{cases} [M]_{0,j} = P_j^{a,b}(-1), \\ [M]_{i,j} = A(x_i) \left[ D_{0,L}^{(a,b,1)} \right]_{i,j} + B(x_i) \left[ {}_C D_{0,L}^{(a,b,\alpha)} \right]_{i,j} + C(x_i) \left[ D_{0,L}^{(a,b,0)} \right]_{i,j}, \\ i = 1, \dots, N, \quad j = 0, 1, \dots, N. \end{cases} \quad (5.11)$$

For the following fractional initial value problem

$$\begin{cases} A(x)u^{(m)}(x) + B(x) {}_C D_{0,x}^\alpha u(x) + C(x)u(x) = f(x), \quad x \in (0, L], \\ u(0) = u_0, \quad u'(0) = u'_0, \end{cases} \quad (5.12)$$

where  $1 < \alpha < 2$ ,  $m = 1$  or  $2$ , we can also get the algebraic equation of the form like (5.10), where  $F = (u_0, f(x_1), f(x_2), \dots, f(x_{N-1}), u'_0)^T$ , and  $M$  satisfies

$$\begin{cases} [M]_{0,j} = P_j^{a,b}(-1), \quad [M]_{N,j} = \frac{2}{L} \frac{dP_j^{a,b}(-1)}{dx}, \\ [M]_{i,j} = A(x_i) \left[ D_{0,L}^{(a,b,m)} \right]_{i,j} + B(x_i) \left[ {}_C D_{0,L}^{(a,b,\alpha)} \right]_{i,j} + C(x_i) \left[ D_{0,L}^{(a,b,0)} \right]_{i,j}, \\ i = 1, \dots, N-1, \quad j = 0, 1, \dots, N. \end{cases} \quad (5.13)$$

If the condition  $u'(0) = u'_0$  in (5.12) is replaced by  $u(L) = u_L$ , then the original fractional initial value problem is reduced to a fractional boundary value problem. We can still get the same form of algebraic equation (5.10), where  $F = (u_0, f(x_1), f(x_2), \dots, f(x_{N-1}), u_L)^T$ , and  $M$  is defined the same as (5.13) except that  $[M]_{N,j} = \frac{2}{L} \frac{dP_j^{a,b}(-1)}{dx}$  is replaced by  $[M]_{N,j} = P_j^{a,b}(1)$ .

## 6. Numerical examples

This section provides numerical examples to verify the methods obtained in the preceding sections. The first two examples are used to test the efficiency and accuracy of the formulae (3.9) and (4.2).

**EXAMPLE 6.1.** Let  $u(x) = x^\mu$ ,  $x \in [0, L] = [0, 1]$ . Now we numerically calculate the fractional integral  $D_{0,x}^{-\alpha} u(x)$  and the Caputo derivative  ${}_C D_{0,x}^\alpha u(x)$ ,  $\alpha > 0$ .

The analytical forms of the fractional integral and the Caputo derivative of  $u(x)$  are given by

$$D_{0,x}^{-\alpha} x^\mu = \frac{\Gamma(\mu+1)}{\Gamma(\mu+1+\alpha)} x^{\mu+\alpha}, \quad \mu > -1,$$

TABLE 1. The absolute errors for Example 6.1 with  $a = b = 0$  and  $\mu = 3.5$ .

| $N$ | $\alpha = 0.2$ | $\alpha = 0.5$ | $\alpha = 0.8$ | $\alpha = 1.2$ | $\alpha = 1.5$ | $\alpha = 1.8$ |
|-----|----------------|----------------|----------------|----------------|----------------|----------------|
| 10  | 4.57e-08       | 3.57e-08       | 1.78e-08       | 5.18e-09       | 1.67e-09       | 6.04e-10       |
| 20  | 2.89e-10       | 1.52e-10       | 5.37e-11       | 9.88e-12       | 2.54e-12       | 1.31e-12       |
| 40  | 1.82e-12       | 6.36e-13       | 1.52e-13       | 1.74e-14       | 3.68e-15       | 2.77e-15       |
| 80  | 1.12e-14       | 2.59e-15       | 4.11e-16       | 1.67e-16       | 1.67e-16       | 1.18e-16       |
| 10  | 2.32e-07       | 1.82e-06       | 8.40e-06       | 1.61e-04       | 3.14e-04       | 3.18e-04       |
| 20  | 2.49e-09       | 2.90e-08       | 1.99e-07       | 6.63e-06       | 1.92e-05       | 2.67e-05       |
| 40  | 2.70e-11       | 4.73e-10       | 4.88e-09       | 2.81e-07       | 1.22e-06       | 2.55e-06       |
| 80  | 2.88e-13       | 7.62e-12       | 1.19e-10       | 1.18e-08       | 7.77e-08       | 2.44e-07       |

and

$${}_CD_{0,x}^\alpha x^\mu = \frac{\Gamma(\mu+1)}{\Gamma(\mu+1-\alpha)} x^{\mu-\alpha}, \quad \mu > -1.$$

Suppose that  $p_N(x)$  is the interpolation of  $u(x)$  on the Jacobi–Gauss–Lobatto points  $x_j$  ( $j = 0, 1, \dots, N$ ) on the interval  $[0, L]$ , and  $p_N$  is expressed by

$$p_N(x) = \sum_{j=0}^N \tilde{p}_j^{a,b} P_j^{a,b} \left( \frac{2x}{L} - 1 \right), \quad (6.1)$$

where  $\tilde{p}_j^{a,b}$  can be easily calculated like (3.2).

We first set  $a = b = 0$  and choose different  $N$  and  $\alpha$  for our computation, the results are shown in Table 1. The first four rows of Table 1 show the absolute maximum errors at the Legendre–Gauss–Lobatto points by using the formula (5.4) (or (3.9)) for the fractional integral. The last four rows of Table 1 give the absolute maximum errors at the Legendre–Gauss–Lobatto points by using the formula (5.5) for the Caputo derivative. Obviously, Table 1 displays the spectral accuracy of the derived method for the approximation of the fractional integral and Caputo derivative.

Now we set  $a = b = -1/2$  to test our algorithms. The numerical results are shown in Table 2. The first four rows and the last four rows give the maximum errors for numerical solutions of the fractional integral and the Caputo derivative of  $u(x)$  at the Chebyshev–Gauss–Lobatto points. We can see that the spectral accuracy is achieved.

**EXAMPLE 6.2.** Let  $u(x) = \sin x$ ,  $x \in [0, L]$ , and we calculate the fractional integral  $D_{0,x}^{-\alpha} u(x)$  and  ${}_CD_{0,x}^\alpha u(x)$ ,  $\alpha > 0$ .

TABLE 2. The absolute errors for Example 6.1 with  $a = b = -1/2$  and  $\mu = 3.5$ .

| $N$ | $\alpha = 0.2$ | $\alpha = 0.5$ | $\alpha = 0.8$ | $\alpha = 1.2$ | $\alpha = 1.5$ | $\alpha = 1.8$ |
|-----|----------------|----------------|----------------|----------------|----------------|----------------|
| 10  | 5.49e-08       | 4.59e-08       | 2.54e-08       | 8.33e-09       | 3.09e-09       | 1.67e-09       |
| 20  | 3.08e-10       | 1.96e-10       | 7.62e-11       | 1.70e-11       | 4.97e-12       | 2.93e-12       |
| 40  | 1.81e-12       | 7.79e-13       | 2.14e-13       | 3.23e-14       | 8.09e-15       | 5.61e-15       |
| 80  | 1.06e-14       | 3.05e-15       | 5.66e-16       | 3.33e-16       | 1.80e-16       | 1.73e-16       |
| 10  | 2.23e-07       | 1.48e-06       | 7.44e-06       | 1.40e-04       | 3.23e-04       | 4.55e-04       |
| 20  | 2.12e-09       | 2.11e-08       | 1.62e-07       | 5.37e-06       | 1.86e-05       | 3.95e-05       |
| 40  | 2.15e-11       | 3.22e-10       | 3.77e-09       | 2.17e-07       | 1.14e-06       | 3.66e-06       |
| 80  | 2.20e-13       | 5.00e-12       | 8.89e-11       | 8.92e-09       | 7.10e-08       | 3.45e-07       |

The exact expressions of  $D_{0,x}^{-\alpha} \sin x$  and  ${}_CD_{0,x}^{\alpha} \sin x$  are given by

$$D_{0,x}^{-\alpha} \sin x = x^{\alpha} \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k+1}}{\Gamma(\alpha + 2k + 1)}, \quad \alpha > 0,$$

and

$$\begin{cases} {}_CD_{0,x}^{\alpha} \sin(x) = x^{1-\alpha} \sum_{k=0}^{\infty} \frac{(-1)^k x^{2k}}{\Gamma(2k + 2 - \alpha)}, & 0 < \alpha < 1, \\ {}_CD_{0,x}^{\alpha} \sin(x) = x^{3-\alpha} \sum_{k=0}^{\infty} \frac{(-1)^{k+1} x^{2k}}{\Gamma(2k + 4 - \alpha)} & 1 < \alpha < 2. \end{cases}$$

We use formula (6.1) as that used in Example 6.1. The numerical experiments are displayed in Tables 3 and 4. The first three rows and the last three rows of Table 3 display the absolute maximum errors of the numerical solutions of the fractional integral and the Caputo derivative of  $u(x)$  by using the methods (5.4) and (5.5) with  $a = b = 0$ , respectively. Table 4 gives the corresponding errors of the case  $a = b = -1/2$ . We can see that the satisfactory results are obtained.

Next, we use our method to solve some fractional differential equations.

EXAMPLE 6.3. Consider the following Bagley–Torvik equation [17, 6]

$$u''(x) + {}_CD_{0,x}^{1.5} u(x) + u(x) = f(x), \quad x \in (0, 1], \quad (6.2)$$

with the initial conditions

$$u(0) = 0, \quad u'(0) = w. \quad (6.3)$$



TABLE 3. The absolute errors for Example 6.2 with  $a = b = 0$ .

| $N$ | $\alpha = 0.2$ | $\alpha = 0.5$ | $\alpha = 0.8$ | $\alpha = 1.2$ | $\alpha = 1.5$ | $\alpha = 1.8$ |
|-----|----------------|----------------|----------------|----------------|----------------|----------------|
| 4   | 4.46e-06       | 5.40e-06       | 3.88e-06       | 1.71e-06       | 6.94e-07       | 2.90e-07       |
| 8   | 4.79e-12       | 4.72e-12       | 2.73e-12       | 8.94e-13       | 2.88e-13       | 6.96e-14       |
| 16  | 6.66e-16       | 2.22e-16       | 2.22e-16       | 1.67e-16       | 1.67e-16       | 8.33e-17       |
| 4   | 1.05e-05       | 6.17e-05       | 2.31e-04       | 1.02e-03       | 2.18e-03       | 5.66e-03       |
| 8   | 1.48e-11       | 1.08e-10       | 5.96e-10       | 4.14e-09       | 1.47e-08       | 5.33e-08       |
| 16  | 3.22e-15       | 1.91e-14       | 9.29e-14       | 6.39e-13       | 2.45e-12       | 9.03e-12       |

TABLE 4. The absolute errors for Example 6.2 with  $a = b = -1/2$ .

| $N$ | $\alpha = 0.2$ | $\alpha = 0.5$ | $\alpha = 0.8$ | $\alpha = 1.2$ | $\alpha = 1.5$ | $\alpha = 1.8$ |
|-----|----------------|----------------|----------------|----------------|----------------|----------------|
| 4   | 6.08e-06       | 7.91e-06       | 6.26e-06       | 3.41e-06       | 1.87e-06       | 1.27e-06       |
| 8   | 6.58e-12       | 6.63e-12       | 3.96e-12       | 1.38e-12       | 4.97e-13       | 1.36e-13       |
| 16  | 3.33e-16       | 3.33e-16       | 2.22e-16       | 5.55e-17       | 2.22e-16       | 5.55e-17       |
| 4   | 1.34e-05       | 5.66e-05       | 1.95e-04       | 9.93e-04       | 2.05e-03       | 5.40e-03       |
| 8   | 1.98e-11       | 1.04e-10       | 3.92e-10       | 3.34e-09       | 1.15e-08       | 4.34e-08       |
| 16  | 4.44e-16       | 1.22e-15       | 7.55e-15       | 4.40e-14       | 2.32e-13       | 9.82e-13       |

Choosing appropriate  $f(x)$  such that (6.2) has the exact solution  $u(x) = \sin wx$ .

We first set  $a = b = 0$ , and the results are shown in Table 5, which shows the maximum absolute errors of the method (5.13) and the shifted Chebyshev tau (SCT) method developed in [6] for the same parameters. From this example, we can see that our method gives the more accurate results than the SCT method used in [6]. Table 6 gives the maximum absolute errors for (6.2) with the boundary value conditions, say, the condition  $u'(0) = w$  is replaced by  $u(1) = \sin w$ . From Tables 5 and 6, we find that the spectral accuracy is attained.

Tables 7 and 8 show the maximum absolute errors of the method (5.13) with  $a = b = -1/2$ . Table 7 displays the errors of the numerical solutions of (6.2) with the initial conditions (6.3), and Table 8 gives the errors of the numerical solutions of (6.2) with the boundary value conditions as mentioned above. Both two cases in this example give the better results than the SCT method reported in [6].

This equation is often used to test the numerical algorithms. In the following we consider the homogeneous Baglay–Torvik model.

TABLE 5. The absolute errors for Example **6.3** with different  $N$  and  $a = b = 0$ .

| $N$ | $w$ | Method (5.13) | SCT [6] | $w$    | Method (5.13) | SCT [6] |
|-----|-----|---------------|---------|--------|---------------|---------|
| 4   | 1   | 2.42e-04      | 3.4e-04 | $4\pi$ | 1.25e+01      | 3.9e-00 |
| 8   | 1   | 7.40e-10      | 4.3e-07 | $4\pi$ | 1.38e+00      | 4.7e-01 |
| 16  | 1   | 3.33e-16      | 1.8e-08 | $4\pi$ | 8.55e-05      | 3.5e-05 |
| 32  | 1   | 4.44e-16      | 7.1e-10 | $4\pi$ | 5.10e-13      | 1.4e-06 |
| 48  | 1   | 3.33e-16      | 9.9e-11 | $4\pi$ | 6.81e-13      | 1.9e-07 |
| 64  | 1   | 4.44e-16      | 2.4e-11 | $4\pi$ | 1.90e-13      | 4.8e-08 |

TABLE 6. The absolute errors for Example **6.3** with boundary value conditions and  $a = b = 0$ .

| $N$ | $w$ | error    | $w$    | error    | $w$    | error    |
|-----|-----|----------|--------|----------|--------|----------|
| 4   | 1   | 2.39e-05 | $2\pi$ | 1.51e-01 | $4\pi$ | 3.47e+00 |
| 8   | 1   | 7.53e-11 | $2\pi$ | 9.20e-04 | $4\pi$ | 1.62e-01 |
| 16  | 1   | 1.11e-16 | $2\pi$ | 1.07e-10 | $4\pi$ | 8.51e-06 |
| 32  | 1   | 3.33e-16 | $2\pi$ | 1.78e-15 | $4\pi$ | 1.13e-12 |
| 48  | 1   | 2.22e-16 | $2\pi$ | 1.47e-15 | $4\pi$ | 2.89e-13 |
| 64  | 1   | 2.22e-16 | $2\pi$ | 1.44e-15 | $4\pi$ | 1.22e-13 |

TABLE 7. The absolute errors for Example **6.3** with initial conditions and  $a = b = -1/2$ .

| $N$ | $w$ | error    | $w$    | error    | $w$    | error    |
|-----|-----|----------|--------|----------|--------|----------|
| 4   | 1   | 1.41e-04 | $2\pi$ | 4.59e-01 | $4\pi$ | 1.36e+01 |
| 8   | 1   | 2.89e-10 | $2\pi$ | 3.40e-03 | $4\pi$ | 5.28e-01 |
| 16  | 1   | 5.55e-16 | $2\pi$ | 2.96e-10 | $4\pi$ | 2.33e-05 |
| 32  | 1   | 4.44e-16 | $2\pi$ | 2.00e-15 | $4\pi$ | 3.32e-13 |
| 48  | 1   | 4.44e-16 | $2\pi$ | 5.33e-15 | $4\pi$ | 3.70e-13 |
| 64  | 1   | 5.55e-16 | $2\pi$ | 3.87e-15 | $4\pi$ | 2.85e-13 |


EXAMPLE **6.4**. We consider the homogeneous Baglay–Torvik equation with order  $\alpha \in (1, 2)$  [1, 17, 7]

$$u''(x) + {}_CD_{0,t}^\alpha u(x) + u(x) = 0, \quad x \in (0, L], \quad u(0) = 1, \quad u'(0) = 0. \quad (6.4)$$

We apply the method (5.13) with  $a = b = 0$  to solve the equation (6.4), where the collocation points are chosen as the Legendre–Gauss–Lobatto points. We set  $L = 50, N = 256$  for different  $\alpha$  in Figures 3 and 4. When

TABLE 8. The absolute errors for Example **6.3** with boundary value conditions and  $a = b = -1/2$ .

| $N$ | $w$ | error    | $w$    | error    | $w$    | error    |
|-----|-----|----------|--------|----------|--------|----------|
| 4   | 1   | 4.43e-05 | $2\pi$ | 3.94e-01 | $4\pi$ | 6.51e+00 |
| 8   | 1   | 6.66e-11 | $2\pi$ | 8.71e-04 | $4\pi$ | 1.97e-01 |
| 16  | 1   | 5.55e-16 | $2\pi$ | 5.04e-11 | $4\pi$ | 4.19e-06 |
| 32  | 1   | 5.55e-16 | $2\pi$ | 1.61e-15 | $4\pi$ | 3.21e-13 |
| 48  | 1   | 5.55e-16 | $2\pi$ | 8.88e-16 | $4\pi$ | 4.45e-13 |
| 64  | 1   | 5.55e-16 | $2\pi$ | 1.39e-15 | $4\pi$ | 3.53e-13 |

FIGURE 3. Numerical result for Example **6.4** with  $N = 256$ ,  $L = 40$ .

$\alpha = 1.25$ , our result coincides with the results reported in [7], see Figure 3. Figure 4 is also consistent with the numerical result in [7].

At last, we use scheme (5.13) to solve an oscillation model.

EXAMPLE **6.5**. Consider the following fractional oscillation equation [7]

$${}_CD_{0,t}^\alpha u(x) + u(x) = f(x), \quad 1 < \alpha < 2, \quad x \in (0, L], \quad (6.5)$$

with the initial conditions

$$u(0) = 1, \quad u'(0) = 0.$$




FIGURE 4. Numerical results for Example 6.4 with  $N = 256$ ,  $L = 40$  and  $\alpha = 1.5, 1.75$ .




FIGURE 5. Numerical results for Example 6.5 with  $N = 256$ ,  $L = 50$ .

In this example, we still use the method (5.13) with  $a = b = 0$  to solve (6.5), where the Legendre–Gauss–Lobatto collocation points are used again. We still let  $N = 256$ ,  $L = 50$  for our computation, and set  $\alpha = 1.3, 1.5, 1.8, 1.95$  as in [7]. The numerical result is displayed in Figure 5, which coincides with the result in [7].

## 7. Conclusion

In this paper, based on the Legendre, Chebyshev and Jacobi polynomials we develop the effective numerical algorithms to compute the fractional integral and Caputo derivative. The operational differential matrix based on the Jacobi–Gauss–Lobatto points is also obtained, which can be seen as a generalization of the classical differential matrix. The computational cost for deriving the operational differential matrix is  $O(N^2)$ , which is much less than  $O(N^3)$  in [5, 6].

## Acknowledgements

The present work was supported by the National Natural Science Foundation of China (Grant no. 10872119), the Shanghai Leading Academic Discipline Project (Grant no. S30104), and the Key Program of Shanghai Municipal Education Commission (Grant no. 12ZZ084).

## References

- [1] R.L. Bagley, P.J. Torvik, On the appearance of the fractional derivative in the behavior of real materials. *J. Appl. Mech.* **51** (1984), 294–298.
- [2] C. Canuto, M.Y. Hussaini, A. Quarteroni, T.A. Zang, *Spectral Methods. Fundamentals in Single Domains*. Springer-Verlag, Berlin (2006).
- [3] F. Cortés, M. Elejabarrieta, Finite element formulations for transient dynamic analysis in structural systems with viscoelastic treatments containing fractional derivative models. *Int. J. Numer. Meth. Engng.* **69** (2007), 2173–2195.
- [4] K. Diethelm, N.J. Ford, A.D. Freed, Y. Luchko, Algorithms for the fractional calculus: a selection of numerical methods. *Comput. Methods Appl. Mech. Engrg.* **194** (2005), 743–773.
- [5] E.H. Doha, A.H. Bhrawy, S.S. Ezz-Eldien, A chebyshev spectral method based on operational matrix for initial and boundary value problems of fractional order. *Comput. Math. Appl.* **62** (2011), 2364–2373.
- [6] E.H. Doha, A.H. Bhrawy, S.S. Ezz-Eldien, Efficient Chebyshev spectral methods for solving multi-term fractional orders differential equations. *Appl. Math. Modelling* **35** (2011), 5662–5672.

- [7] J.T. Edwards, N.J. Ford, A.C. Simpson, The numerical solution of linear multi-term fractional equations: System of equations. *J. Comput. Appl. Math.* **148** (2002), 401–418.
- [8] S. Esmaeili, M. Shamsi, A pseudo-spectral scheme for the approximate solution of a family of fractional differential equations. *Commun. Nonlinear Sci. Numer. Simulat.* **16** (2011), 3646–3654.
- [9] V.V. Kulish, Application of fractional calculus to fluid mechanics. *J. Fluids Eng.* **124** (2002), 803–808.
- [10] C.P. Li, A. Chen, J.J. Ye, Numerical approach to fractional calculus and fractional ordinary differential equations. *J. Comput. Phys.* **230** (2011), 3352–3368.
- [11] C.P. Li, F.H. Zeng, Finite difference methods for fractional differential equations. *Int. J. Bifurcation Chaos* **22**, No 4 (2012), 1230014 (28 pages); DOI: 10.1142/S0218127412300145; <http://www.worldscinet.com/ijbc/ijbc.shtml>
- [12] F. Liu, Q.Q. Yang, I. Turner, Two new implicit numerical methods for the fractional cable equation. *J. Comput. Nonlinear Dyn.* **6** (2011), 011009–1.
- [13] C. Lubich, Discretized fractional calculus. *SIAM J. Math. Anal.* **17** (1986), 704–719.
- [14] V.E. Lynch, B.A. Carreras, D. del-Castillo-Negrete, K.M. Ferreira-Mejias, H.R. Hicks, Numerical methods for the solution of partial differential equations of fractional order. *J. Comput. Phys.* **192** (2003), 406–421.
- [15] K. Miller, B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*. Wiley, New York (1993).
- [16] K.B. Oldham, J. Spanier, *The Fractional Calculus*. Academic Press, New York (2006).
- [17] I. Podlubny, *Fractional Differential Equations*. Academic Press, San Diego (1999).
- [18] I. Podlubny, Matrix approach to discrete fractional calculus, *Fract. Calc. Appl. Anal.* **3**, No 4 (2000), 359–386.
- [19] I. Podlubny, A. Chechkin, T. Skovranek, Y.Q. Chen, B. Vinagre, Matrix approach to discrete fractional calculus II: Partial fractional differential equations. *J. Comput. Phys.* **228** (2009), 3137–3153.

- [20] Y.A. Rossikhin, M.V. Shitikova, Application of fractional calculus for dynamic problems of solid mechanics: Novel trends and recent results. *Applied Mechanics Reviews* **63** (2010), 010801–1.
- [21] A. Saadatmandi, M. Dehghan, A new operational matrix for solving fractional-order differential equations. *Comput. Math. Appl.* **59** (2010), 1326–1336.
- [22] S.G. Samko, A.A. Kilbas, O.I. Marichev, *Fractional Integrals and Derivatives*. Gordon and Breach Science, Yverdon, Switzerland (1993).
- [23] A. Schmidt, L. Gaul, On the numerical evaluation of fractional derivatives in multi-degree-of-freedom systems. *Signal Processing* **86** (2006), 2592–2601.
- [24] J. Shen, T. Tang, L. L. Wang, *Spectral Methods. Algorithms, Analysis and Applications*. Springer-Verlag, Heidelberg, Berlin (2011).
- [25] E. Sousa, How to approximate the fractional derivative of order  $1 < \alpha \leq 2$ . In: *Proceedings of FDA'10. The 4th IFAC Workshop Fractional Differentiation and its Applications*, Badajoz, Spain (2010).
- [26] H. Sugiura, T. Hasegawa, Quadrature rule for Abel's equations: Uniformly approximating fractional derivatives. *J. Comput. Appl. Math.* **223** (2009), 459–468.
- [27] J. Tenreiro Machado, V. Kiryakova, F. Mainardi, Recent history of fractional calculus. *Commun. Nonlinear Sci. Numer. Simul.* **16**, No 3 (2011), 1140–1153; doi:10.1016/j.cnsns.2010.05.027; <http://www.sciencedirect.com/science/article/pii/S1007570410003205>
- [28] H.G. Sun, W. Chen, C.P. Li, Y.Q. Chen, Finite difference schemes for variable-order time fractional diffusion equation. *Int. J. Bifurcation Chaos* **22**, No 4 (2012), 1250085 (16 pages); DOI: 10.1142/S021812741250085X; <http://www.worldscinet.com/ijbc/ijbc.shtml>
- [29] Z.Z. Sun, X.N. Wu, A fully discrete difference scheme for a diffusion-wave system. *Appl. Numer. Math.* **56** (2006), 193–209.
- [30] Q. Yu, F. Liu, V. Anh, I. Turner, Solving linear and nonlinear space-time fractional reaction-diffusion equations by the Adomian decomposition method. *Int. J. Numer. Meth. Engng.* **74** (2008), 138–158.
- [31] Z.M. Odibat, Computational algorithms for computing the fractional derivatives of functions. *Math. Comput. Simulat.* **79** (2009), 2013–2020.
- [32] P. Zhuang, T. Gu, F. Liu, I. Turner, P.K.D.V. Yarlagadda, Time-dependent fractional advection-diffusion equations by an implicit MLS meshless method. *Int. J. Numer. Meth. Engng.* **88** (2011), 1346–1362.

<sup>1,2</sup> *Department of Mathematics*  
*Shanghai University*  
*Shanghai – 200444, PR CHINA*  
*e-mail: lcp@shu.edu.cn*

*Received: February 11, 2012*

<sup>3</sup> *School of Mathematical Sciences*  
*Queensland University of Technology*  
*GPO Box 2434, Brisbane, Qld.*  
*4001, AUSTRALIA*  
*e-mail: f.liu@qut.edu.au*

---

Please cite to this paper as published in:

*Fract. Calc. Appl. Anal.*, Vol. **15**, No 3 (2012), pp. 383–406;  
DOI: 10.2478/s13540-012-0028-x