CME quiz

The purpose of this quiz is to provide a convenient means of self-assessment of your reading of the scientific content of this issue of *JAOA*. Enter your answers to the questions in the spaces provided so that you can easily check them with the answers that will be published next month.

To apply for CME credit, transfer your answers to the application form and answer sheet on page 318 and mail it to the CME office. So that you may complete this self-assessment in privacy, use only your member number to apply for CME credit. The CME office will record only the fact that you have completed the self-assessment test. Any grading will be done by the Editorial Department only for the purpose of planning areas of study which may be helpful to cover in future issues of *JAOA*.

- 1. All the following osteopathic manipulative techniques are indicated for the treatment of juvenile rheumatoid arthritis (JRA) except:
- ____(a) High-velocity/low-amplitude techniques.
- —(b) Lymphatic pump.
- ___(c) Joint pump.
- ___(d) Jones' technique.
- ——(e) Myofascial techniques. 2. Which of the following JRA su
- 2. Which of the following JRA subtypes/subsets is most closely associated with the loss of visual acuity?
- ____(a) Polyarticular/rheumatoid factor-positive.
- ____(b) Polyarticular/rheumatoid factor-negative.
- ——(c) Pauciarticular/antinuclear antibody—positive.
- ——(d) Pauciarticular/antinuclear antibody–negative.
- ——(e) Pauciarticular/histocompatibility locus antigen B-27– positive.
- 3. Which of the following statements are true concerning preemptive analgesia?

- ____(a) It may prevent peripheral sensitization.
- ____(b) It may prevent central sensitization.
- (c) It may be useful in attenuating or preventing postoperative pain.
- ___(d) All the above.
- 4. Which of the following statements are true concerning the American Academy of Pediatrics guidelines during medical procedures in children?
- (a) During conscious sedation, the child is able to maintain a patent airway independently and continuously.
- (b) During deep sedation, the child may not be able to independently maintain his or her airway.
- ——(c) Continuous pulse oximetry and heart rate monitoring are required during conscious sedation.
- ——(d) All the above.
- 5. The tissue diagnosis most commonly found with a giant sigmoid diverticulum is which of the following?
- ___(a) Pseudocyst.
 - __(b) True diverticulum.
- ____(c) Adenocarcinoma of the wall.
- ___(d) None of the above.
- 6. Peritonitis is the most frequent finding at the initial examination.
- ___(a) True.
 - __(b) False.
- 7. The most common site of metastasis of renal cell carcinoma is which of the following?
 - __(a) Bone.
- ___(b) Lung.
 - __(c) Pancreas.
- ___(d) Regional lymph nodes.
- —(e) Vena cava.
- 8. The most common adjacent structure to be involved with metastatic renal cell carcinoma is which of the following?
- ___(a) Brain.
 - _(b) Colon.
- ___(c) Pancreas.
 - ___(d) Regional nodes.
- ____(e) Contralateral kidney.

advertisers' index

American Osteopathic Association

CME Video: "Back to the Future: Primary Care in America," 265

Forest Laboratories

Esgic Plus, 281, 292

Janssen Pharmaceutica

Propulsid, 269, 270 Sporanox, 277, 278

Procter & Gamble

Olean, 263

Westview Hospital, 317

Wyeth-Ayerst Laboratories

Lodine, Cover 2, 259, 260

Zeneca Pharmaceuticals

Sular, 267, 268, 303, 304 Zestril, Cover 3, Cover 4