

The purpose of this quiz is to provide a convenient means of self-assessment of your reading of the scientific content of this issue of *JAOA*. Enter your answers to the questions in the spaces provided so that you can easily check them with the answers that will be published next month.

To apply for CME credit, transfer your answers to the application form and answer sheet on page 375 and mail it to the CME office. So that you may complete this self-assessment in privacy, use only your member number to apply for CME credit. The CME office will record only the fact that you have completed the self-assessment test. Any grading will be done by the Editorial Department only for the purpose of planning areas of study which may be helpful to cover in future issues of *JAOA*.

1. The risk of adverse effects to astemizole is increased with all the following *except*:

- _____ (a) Erythromycin.
- _____ (b) Terfenadine.
- _____ (c) Loading dose.
- _____ (d) Prednisone.
- _____ (e) Ketoconazole.

2. All the following may be considered etiologies of a spontaneous hemoperitoneum *except*:

- _____ (a) Ruptured abdominal aortic aneurysm.
- _____ (b) Ruptured ovarian cyst.
- _____ (c) Portal hypertension.
- _____ (d) Pancreatitis.
- _____ (e) Vasculitis.

3. Massive hemoperitoneum is always preceded by an inciting event.

- _____ (a) True.
- _____ (b) False.

4. Characteristic features of Sweet's syndrome include all the following *except*:

- _____ (a) Tender erythematous or violaceous cutaneous plaques or nodules of abrupt onset.
- _____ (b) Vasculitis.
- _____ (c) Preceding fever or infection.
- _____ (d) Accompanying fever, arthralgia, conjunctivitis, or malignancy.

_____ (e) Erythrocyte sedimentation rate greater than 50 mm/h.

5. What percentage of cases of malignancy-associated Sweet's syndrome are hematologic?

- _____ (a) Twenty percent.
- _____ (b) Fifty-five percent.
- _____ (c) Seventy percent.
- _____ (d) Eighty-five percent.
- _____ (e) None of the above.

6. What is the minimum latency period required for a radiation-induced malignancy to develop?

- _____ (a) Two months.
- _____ (b) Two years.
- _____ (c) Five years.
- _____ (d) Fifteen years. ♦

Build Your Family Practice.

Start with a solid foundation at the center of osteopathic medicine for central Indiana. Add to this our exceptional professional opportunities, competitive salary and excellent benefits package.

You'll be a member of the prestigious medical community of Indianapolis, the nation's 12th largest city with many opportunities for growth.

Join the Westview team. And see what develops for your private practice.

Westview Hospital

AN OSTEOPATHIC INSTITUTION

3630 Guion Road
Indianapolis, Indiana 46222

For more information contact:
Michele Millikan at 1-800-587-7085.